

Self-properties of codimension-2 braneworlds

Christos Charmousis^a and Antonios Papazoglou^b

^a*LPT, Université de Paris-Sud,
Bât. 210, 91405 Orsay CEDEX, France*

^b*Institute of Cosmology and Gravitation,
University of Portsmouth, Portsmouth PO1 2EG, U.K.*

E-mail: christos.charmousis@th.u-psud.fr, antonios.papazoglou@port.ac.uk

ABSTRACT: We consider four-dimensional de Sitter, flat and anti de Sitter branes embedded in a six-dimensional bulk spacetime whose dynamics is dictated by Lovelock theory. We find, applying a generalised version of Birkhoff's theorem, that all possible maximally symmetric braneworld solutions are embedded in Wick-rotated black hole spacetimes of Lovelock theory. These are warped solitonic spaces, where the horizons of the black hole geometries correspond to the possible positions of codimension-2 branes. The horizon temperature is related via conical singularities to the tension or vacuum energy of the branes. We classify the braneworld solutions for certain combinations of bulk parameters, according to their induced curvature, their vacuum energy and their effective compactness in the extra dimensions. The bulk Lovelock theory gives an induced gravity term on the brane, which, we argue, generates four-dimensional gravity up to some distance scale. As a result, some simple solutions, such as the Lovelock corrected Schwarzschild black hole in six dimensions, are shown to give rise to self-accelerating braneworlds. We also find that several other solutions have self-tuning properties. Finally, we present regular gravitational instantons of Lovelock gravity and comment on their significance.

KEYWORDS: p-branes, Large Extra Dimensions, Solitons Monopoles and Instantons, Black Holes.

Contents

1. Introduction	1
2. Static black hole solutions	4
3. Codimension-2 braneworlds	7
4. Braneworld solutions	9
4.1 Zero bare cosmological constant	9
4.1.1 The dS_4 vacua	10
4.1.2 The AdS_4 vacua	11
4.1.3 The Nariai dS_4 vacuum	11
4.1.4 Zero black hole mass $\mu = 0$	12
4.1.5 Flat vacuum	13
4.2 Non-zero bulk cosmological constant	13
4.2.1 Zero black hole mass $\mu = 0$	13
4.2.2 The Born-Infeld vacua	14
4.2.3 The Born-Infeld Nariai vacuum	15
4.2.4 Flat vacua	16
5. Self-properties of the solutions	16
6. The Gauss-Bonnet instantons	20
7. Discussion and conclusions	21
A. A more general action	24
B. The general solution of $x^3 + Ax + M = 0$	25
C. Volume calculation	26

1. Introduction

Combined cosmological and astrophysical data indicate that at least 96% of the actual matter content of the Universe has yet to be detected in particle accelerators. This is a correct statement provided we assume a homogeneous universe described by Einstein's field equations. One fourth of this yet unseen component, dark matter, we can hope to discover in the Large Hadron Collider and is an expected, quite natural, cosmological signature of particle physics theories such as low energy supersymmetry (SUSY). For the

remaining chunk, usually dubbed dark energy, that has cropped up more recently from supernovae data [1], the situation is a lot less clearer. Data is best "fitted" by assuming a tiny positive cosmological constant of magnitude $(10^{-3}\text{eV})^4$. The mass scale associated to this energy is that of, roughly, what one expects for neutrino masses, a really tiny number in the cosmological arena. This cosmological constant is so small because the radius of the observed universe is by now huge, of size of the square inverse to the mass scale in question. It is dominant because dark matter is extremely diluted at cosmological scales of roughly 10Gpc whereas super clusters of galaxies such as Virgo are at 20Mpc.

Unlike dark matter, from the point of view of theoretical particle physics, the cosmological constant is rather problematic. For the "natural" value of the cosmological constant, from the point of view of Quantum Field Theory, is of the order of the ultraviolet cut-off we would impose for our quantum field theory, ranging from the Planck scale to the SUSY breaking scale, depending on our theoretical prejudice. The fine tuning we impose to compensate this vacuum energy by some bare, gravitational cosmological constant is the biggest known discrepancy between theoretical expectations and our understanding of experimental data. This is the cosmological constant problem [2] that was there already there [3] when the precision of existing data pointed towards a value, assumed for simplicity to be zero.

By admitting the presence of a small, but not null, cosmological constant we add two additional problems to the "old" cosmological constant problem [3], namely, why is it so small and not zero, and why is its density now of the same order as dark matter density. Why therefore can we observe it now, *i.e.* the right time that it is possible for us to observe it in the first place? Dark energy models usually accompanied by scalar fields with suitable potentials do not solve this fine-tuning problem. They are also generically problematic due to enhanced radiative corrections associated to the scalar(s) [4] coupling to matter. Although general relativity is very well tested at solar system scales, both at weak and strong field regimes [5], it is not so at cosmological scales. In fact, to get an idea of the colossal difference in scales, take the quotient of the Hubble radius over the Astronomical unit (earth to sun distance) to get 10^{15} ! Therefore, cosmological data, if we do not assume some extra source of unknown matter or cosmological constant, tell us precisely that gravity is modified at the infrared.

Taking into account the above discussion, modifying gravity in the infrared is a legitimate theoretical hypothesis that should be taken seriously. As yet, no convincing modification of general relativity has been found. Often the problem is experimental, constraints as for example for $f(R)$ theories in the solar system and in galaxy clusters [6]. However, there are also theoretical difficulties since a large distance modification spoils our understanding of asymptotic behaviour, which is very well-defined in General Relativity (GR). Typically, such models [7, 8] have ghost instabilities [9–11] and strong coupling problems [12] which are as yet not understood and therefore are no better than fitting data with a single, however small parameter, the cosmological constant.

This of course does not mean there is no such consistent modification. Clearly one has to try more complex setups and some interesting toy models [13] have been proposed and are now being put to the test. By far, however, the most successful and popular

modification is that of self-acceleration [14] present in the Dvali-Gabadagze-Porrati (DGP) model [8]. There, the small current acceleration of the Universe is understood as a geometric effect originating from the (unfair) competition between the five-dimensional and four-dimensional curvature scales. Unfair, because the resulting length scale, the crossover scale, is of the order of the Hubble horizon today, $H_0 \sim 10^{-34} eV$. At these length scales gravity becomes five-dimensional with a very low five-dimensional gravitational scale, roughly of $10^4 GeV$ and we enter a geometric acceleration phase. Unfortunately, this solution has a ghost in its linear perturbation spectrum [10, 11]. Furthermore, the results are fuzzed by the presence of strong coupling problems [12] and either the theory loses linear predictability (which makes it useless cosmologically) or is completely unstable (see [15] for exact solutions and beyond linear order). To summarize, therefore, self-acceleration teaches us that we can in principle explain in a simple way the current acceleration of the Universe by a geometric effect of a codimension-1 braneworld, but the geometry modification required to attain this result is not seemingly viable. Along these lines, recently [16] have given an idea on similar terms where the cosmological constant results from a higher order gravity correction in a codimension-1 braneworld.

An even more ambitious proposal that emerged in braneworld models is that of the self-tuning and has to do with the "old" cosmological constant problem [3]. The idea was to find models where the tension of the brane, that is its vacuum energy, can be large without affecting the curvature of the brane and without fine-tuning of it with other brane or bulk parameters [17]. The best attempt to realize such a kind of model has been in six dimensions and in the case that the brane is of codimension-2 [18]. These conical branes have the special property that their tension merely induces a deficit angle in the surrounding bulk geometry and does not curve the brane world-volume. The brane curvature is induced from the bulk dynamics and is not directly related to the brane vacuum energy.¹ Several models with codimension-2 branes were studied with various compactifications [19, 20], however most of them had hidden fine-tunings or curvature singularities present. Although self-tuning vacua can be found, where indeed solutions of the same curvature correspond to a (continuous) range of brane vacuum energy, a cosmological constant problem resolution would require a dynamical selection mechanism of such vacua. This is because, as a rule, there exist nearby solutions of different curvature, which introduce the question whether the self-tuning solution is an attractor.

In the present paper, we will examine a completely novel possibility of obtaining acceleration due to geometry as well as certain self-tuning properties. The modified gravity theory that we will study is Lovelock theory [21] in six dimensions, which is the natural extension of GR in higher dimensions. The Lovelock theory in six dimensions has in addition to the Einstein-Hilbert term the Gauss-Bonnet combination. Although the latter is a topological invariant in four dimensions, it becomes dynamical for higher dimensions and modifies the gravitational theory. The codimension-2 branes present in the generic vacua of this theory, can have interesting properties, similar to the ones stated in the previous para-

¹We emphasize here that we will not necessarily consider flat brane curvature, but also cases where a small brane curvature is allowed with a large vacuum energy on the brane.

graphs. We will find for example self-accelerating cases with non-compact internal spaces as well as self-tuning vacua for several compact and non-compact vacua. These examples open new possibilities for consistent self-acceleration and effective self-tuning which need to be considered in more detail in the future.

In fact, Lovelock theory plays a crucial double role, not only in the bulk, but also for the brane junction conditions [22]. On the one hand, it provides geometric self-acceleration and novel solutions which are absent in Einstein theory, and on the other hand, it gives an induced gravity term on the brane [22, 23], much like in DGP [8]. Here we must emphasise, that this does not mean that we expect to have Einstein gravity on the codimension-2 braneworld, rather, it means that up to some scale, gravity "looks" four-dimensional as in the DGP model. Lovelock theory has the extraordinary geometric property to induce an Einstein-Hilbert term at the level of the junction conditions [23].

The structure of the paper is as follows: we will firstly present the general black hole solutions in six-dimensional Lovelock theory and then show how by double Wick-rotation we can obtain the most general axially symmetric braneworlds with maximally symmetric codimension-2 branes. Then we will scan through several classes of solutions and analyse the cases of interest, *i.e.* the self-accelerating and the self-tuning vacua introducing each time, as few "extra bulk parameters" as possible. We will briefly comment on the physical implications of some new regular instanton solutions of Lovelock gravity and we will finally conclude.

2. Static black hole solutions

Let us consider the six-dimensional dynamics of gravity with a bare cosmological constant Λ and a Gauss-Bonnet (GB) term (see appendix A for a more general system with in addition a gauge field coupled to gravity). The action of the system reads

$$S = \int d^6x \sqrt{-g} \left[\frac{1}{16\pi G_6} (R + \hat{\alpha} \mathcal{L}_{\text{GB}}) - 2\Lambda \right], \quad (2.1)$$

where

$$\mathcal{L}_{\text{GB}} = R_{MNK\Lambda} R^{MNK\Lambda} - 4R_{MN} R^{MN} + R^2, \quad (2.2)$$

is the Gauss-Bonnet Lagrangian density, G_6 the six-dimensional Newton's constant and $\hat{\alpha}$ the Gauss-Bonnet coupling.

It is straightforward to write down the Einstein equations of motion for the above action. They read

$$G_{MN} - \hat{\alpha} H_{MN} = -8\pi G_6 \Lambda g_{MN}, \quad (2.3)$$

with $G_{MN} = R_{MN} - \frac{1}{2} R g_{MN}$ the Einstein tensor and the following Gauss-Bonnet contribution

$$H_{MN} = \frac{1}{2} \mathcal{L}_{\text{GB}} g_{MN} - 2R R_{MN} + 4R_{MK} R_N^K + 4R_{KMAN} R^{K\Lambda} - 2R_{MK\Lambda P} R_M^{K\Lambda P}. \quad (2.4)$$

Our ultimate goal is to find the maximally symmetric solutions of the above equations of motion. For this purpose, let us consider a four-dimensional space of maximal symmetry parametrised by $\kappa = 0, -1, 1$, with metric

$$h_{\mu\nu} = \frac{\delta_{\mu\nu}}{\left(1 + \frac{\kappa}{4}\delta_{\mu\nu}x^\mu x^\nu\right)^2}. \quad (2.5)$$

These spaces are four-dimensional flat space, hyperboloid or sphere respectively of curvature $R[h] = 12\kappa$. A generalised version of Birkhoff's theorem for (2.1) states, [24] (see [25] for the Lovelock version) that every six-dimensional spacetime solution of (2.3) having such four-dimensional maximal sub-spaces (2.5) is locally isometric to

$$ds^2 = -V(r)dt^2 + \frac{dr^2}{V(r)} + r^2 h_{\mu\nu} dx^\mu dx^\nu, \quad (2.6)$$

admitting therefore a locally timelike Killing vector (∂_t in the coordinates of (2.6)). The solution of the potential V for the equations of motion (2.3) is found to be [26, 27]

$$V(r) = \kappa + \frac{r^2}{2\alpha} \left[1 + \epsilon \sqrt{1 + 4\alpha \left(a^2 - \frac{\epsilon\mu}{r^5} \right)} \right], \quad (2.7)$$

where the parameters appearing in the action (2.1) have been rescaled to,

$$\begin{aligned} \alpha &= 6\hat{\alpha}, & 16\pi G_6 \Lambda &= 20a^2 > 0 \quad (\text{for } dS_6), \\ & & a^2 &= -k^2 < 0 \quad (\text{for } AdS_6). \end{aligned} \quad (2.8)$$

The integration constant μ is

$$\mu = \frac{4\pi G_6 M}{\Sigma_\kappa}, \quad (2.9)$$

where M is the AD or ADM mass of the solution and Σ_κ is the area of the unit four-dimensional maximally symmetric subspace. Finally, $\epsilon = \pm 1$, giving rise to two distinct branches of solutions. The convention of the μ sign is chosen so that the gravitational mass is always $\mu > 0$. Indeed, as one can easily check by expanding the square root for large distances, the sign flip in front of μ is necessary to match the Schwarzschild de Sitter solution behaviour for positive AD mass. The case where $1 + 4\alpha a^2 = 0$ is special, because the theory can be written in a Born-Infeld (BI) form [28].

Setting $\mu = 0$ gives us the asymptotic vacua of the theory (2.1) which unlike Einstein theory are not unique (for given bare parameters in (2.1)). Then, we find that asymptotically we have an effective six-dimensional cosmological constant

$$16\pi G_6 \Lambda_{\text{eff}} = -\frac{10}{\alpha} \left[1 + \epsilon \sqrt{1 + 4\alpha a^2} \right], \quad (2.10)$$

with a normalisation like in (2.1). We note in particular, that α gives an effective cosmological constant in the bulk even without a cosmological constant term in the action, *i.e.* even with $a^2 = 0$. The maximally symmetric space that the solutions asymptote to, depends on the sign of α and the branch of the solution, *i.e.* ϵ . It is interesting to expand

the above expression at $\alpha \rightarrow 0$ in order to check if there is an Einstein theory limit for the above vacua or not

$$16\pi G_6 \Lambda_{\text{eff}} = -\frac{10}{\alpha}(1 + \epsilon) - 20\epsilon a^2 + \mathcal{O}(\alpha) . \quad (2.11)$$

Hence, we distinguish the following three cases

- For $\epsilon = +1$, the solution, as can be seen by (2.11), does not have an Einstein theory limit as $\alpha \rightarrow 0$, although there may be a relevant dS_6 or AdS_6 Einstein solution mimicking (2.6). We name this branch the *Gauss-Bonnet branch*. The solution asymptotes AdS_6 space for $\alpha > 0$ and to dS_6 space for $\alpha < 0$. Unlike what is argued in [26] this branch is not, at least, classically unstable for the reasons put forward by [29] (for a full study of this see [30]).
- For $\epsilon = -1$, the solution has an Einstein theory limit (as $\alpha \rightarrow 0$) as seen from (2.11) and therefore we call this branch the *Einstein branch*. The solution asymptotes AdS_6 space for $\alpha > 0$, $-1/(4\alpha) < a^2 < 0$ or $\alpha < 0$, $a^2 < 0$, to dS_6 space for $\alpha < 0$, $0 < a^2 < 1/(4|\alpha|)$ or $\alpha > 0$, $a^2 > 0$ and to M_6 for $a^2 = 0$.
- For the special BI case $1 + 4\alpha a^2 = 0$, as we can see from (2.10) we obtain an dS_6 or AdS_6 asymptotic solution which does not have an Einstein theory limit and no possible flat vacuum. However this is the only case that we have a unique vacuum.

Let us now proceed into analysing the solution at hand (2.6) (we will mostly follow [31]). For the above metric solution (2.6), there are two possible singularities in the curvature tensor, the usual $r = 0$, and also a branch cut singularity at the (maximal) zero of the square root

$$r_s^5 = \frac{4\alpha\epsilon\mu}{1 + 4\alpha a^2} . \quad (2.12)$$

Whenever r_s is real and positive, this is the singular end of spacetime (2.6). For the BI case $1 + 4\alpha a^2 = 0$ there is no such singularity.

We will have a black hole solution if and only if there exists $r = r_h$ such that $V(r_h) = 0$ and $r_h > r_{\text{max}}$, where $r_{\text{max}} = \max\{0, r_s\}$. Indeed the usual Kruskal extension

$$dv_{\pm} = dt \pm \frac{dr}{V(r)} , \quad (2.13)$$

gives that (v_+, r) and (v_-, r) constitute a regular chart across the future and past horizons of (2.6). It is actually straightforward and rather useful to show the following: $r = r_h$ is an horizon iff,

$$r_h > r_{\text{max}} , \quad (2.14)$$

$$\epsilon(2\alpha\kappa + r_h^2) \leq 0 , \quad (2.15)$$

$$p_{\alpha}(r_h) = 0 \quad \text{with } p_{\alpha}(x) = -a^2 x^5 + \kappa x^3 + \alpha \kappa^2 x + \epsilon \mu . \quad (2.16)$$

In addition, one should make sure that the sign of V on each side of the roots of the above polynomial is such that (2.6) describes a black hole. The sign of p_{α} and V are not in general the same, depending on the signs of ϵ and α .

Therefore for $\epsilon = +1$ we have $r_h^2 \leq -2\alpha\kappa$, *i.e.*, the event horizon is bounded from above. This means that for $\kappa = 0$ there are no black hole solutions in this branch. Also note that $p_{\alpha=0}(x)$ is just the usual polynomial for a Einstein black hole, and hence, α couples only to the horizon curvature κ . Therefore, for $\epsilon = -1$ and $\kappa = 0$ the horizon positions are the same as in the GR solutions. In fact, when $\epsilon = -1$ we have similar structure and properties of the solutions (2.6) as their Einstein counterparts.

3. Codimension-2 braneworlds

A standard procedure for Einstein theory to generate brane world solutions from black hole solutions [32, 33] is to perform a double Wick rotation for a black hole solution. The same will be valid for (2.1) and the black hole solution (2.6) at hand. We set $t \rightarrow i\theta$ and in addition make a further Wick rotation $x^0 \rightarrow it$ in the metric $h_{\mu\nu}$ (2.5) so that it becomes of Lorentzian signature

$$h_{\mu\nu} = \frac{\eta_{\mu\nu}}{\left(1 + \frac{\kappa}{4}\eta_{\mu\nu}x^\mu x^\nu\right)^2} . \tag{3.1}$$

Then, the maximally symmetric spacetime sections correspond to four-dimensional Minkowski, AdS_4 and dS_4 for $\kappa = 0, -1, 1$ respectively with curvature $R[h] = 12\kappa$. The solutions are now of manifest axial symmetry with ∂_θ as the angular Killing vector,

$$ds^2 = V(r)d\theta^2 + \frac{dr^2}{V(r)} + r^2 h_{\mu\nu} dx^\mu dx^\nu . \tag{3.2}$$

The six-dimensional spacetime has the correct symmetries to describe a maximally symmetric four-dimensional brane world. The staticity theorem invoked for black hole spacetimes [24] tells us now that axial symmetry comes for free and need not be imposed for resolving the system of equations. The solutions (3.2) are therefore the general solutions describing maximally symmetric branes, something that has been overlooked up to now even in the case of Einstein theory. It is interesting to note that in the case of Einstein theory in four dimensions, we obtain in this way the general maximally symmetric gravitating cosmic strings, which can differ drastically from their flat counterparts (see in particular, the cosmic string solutions in AdS presented in [34]).

The positions of the horizons r_h will be the endpoints of the internal space and the solutions will have meaning if we keep the spacelike regions of the previous black hole solutions, *i.e.* the ones with $V(r) > 0$. Let us suppose that we keep the space-like region between two horizons r_- and r_+ , with $r_- < r_+$ [the subsequent discussion applies of course also in the case that we have only one horizon and we keep the side which is spacelike]. At these endpoints of spacetime, which are also the fixed points of the axial symmetry, one can in general put branes of dimension four, in other words 3-branes. Since the brane solutions we have found are maximally symmetric, the brane can only carry some two-dimensional Dirac charge, *i.e.* pure tension. Taking $x^\mu = \text{const.}$ and expanding around the zeros of V we get,

$$ds^2 \approx \left(\frac{1}{4}V'_{r_\pm}\right) \rho_\pm^2 d\theta^2 + d\rho_\pm^2 , \tag{3.3}$$

with the Gaussian Normal radial coordinate

$$\rho_{\pm} = \sqrt{\frac{4(r - r_{\pm})}{V'_{r_{\pm}}}}, \quad (3.4)$$

which is well defined in all cases with $V'_{r_{\pm}} \neq 0$. The case of $V'_{r_{\pm}} = 0$ needs special attention as we will see later (section 4.1.3 and section 4.2.3). In this coordinate system, the brane energy momentum tensor is $T_{\mu\nu}^{\text{brane}} = S_{\mu\nu}\delta^{(2)} = S_{\mu\nu}\frac{\delta(\rho_{\pm})}{2\pi\rho_{\pm}}$, with $S_{\mu}^{\nu} = -T_{\pm}\delta_{\mu}^{\nu}$, where T_{\pm} are the brane tensions. If the angular coordinate has periodicity $\theta \in [0, 2\pi c)$, then the deficit angles which are induced at the two brane positions are $\delta_{\pm} = 2\pi(1 - \beta_{\pm})$ with

$$\beta_{\pm} = \frac{1}{2}|V'_{\pm}|c. \quad (3.5)$$

The angular periodicity c is an arbitrary topological integration constant of the solution and can be varied to generate physically different brane world solutions. Let us note here, that in a compact model, we can always use the freedom of choosing c to set the deficit angles of one of the branes to zero, *e.g.* $\beta_{-} = 1$, thus, obtaining a "teardrop" compact space with only one brane of $\beta_{+} \neq 1$.

From the Einstein equations (2.3) supplemented by the brane tension terms, one can separate the distributional Dirac parts and write down induced Einstein equations for the branes. These brane junction conditions are [22, 23]

$$2\pi(1 - \beta_{\pm}) \left(-\gamma_{\mu\nu} + 4\hat{\alpha}G_{\mu\nu}^{\text{ind}} \right) = 8\pi G_6 S_{\mu\nu}, \quad (3.6)$$

where $\gamma_{\mu\nu}^{\pm} = r_{\pm}^2 h_{\mu\nu}$ is the induced metric on the branes with curvature $R[\gamma^{\pm}] = 12\kappa/r_{\pm}^2 \equiv 12\kappa H_{\pm}^2$, and $G_{\mu\nu}^{\text{ind}} = -3\kappa H_{\pm}^2 \gamma_{\mu\nu}$ is the induced Einstein tensor. It is important to emphasize here that r_{\pm}^2 depends on the geometric parameters of the bulk solution (3.2) namely the mass μ , and the bare parameters α and a^2 . It is also effectively the warp factor of the brane. The induced Newton's constant on the two branes can be determined from (3.6) to be

$$G_4^{\pm} = \frac{3G_6}{4\pi\alpha(1 - \beta_{\pm})}. \quad (3.7)$$

Note that in order to have positive induced Newton's constant, we should have angle deficit ($\beta_{\pm} < 1$) for $\alpha > 0$ and angle excess ($\beta_{\pm} > 1$) for $\alpha < 0$. The two dimensional warped space in the coordinates θ and r has in the latter case the shape of a pumpkin whereas in the former case the shape of a football (rugby ball to be more precise). The important information coming from (3.6) is that the parameters α and β invoke the relation and the possible hierarchy between the bulk and four-dimensional scales.

Substituting the $G_{\mu\nu}^{\text{ind}}$ back in (3.6) we find a relation between the Hubble parameters on the branes and the action parameters

$$2\pi(1 - \beta_{\pm}) \left(\frac{1}{2\alpha} + \kappa H_{\pm}^2 \right) = \frac{4\pi G_6}{\alpha} T^{\pm}, \quad (3.8)$$

which can be further simplified, if we substitute the deficit angle from (3.7) and solve for H_{\pm}^2

$$\kappa H_{\pm}^2 = -\frac{1}{2\alpha} + \frac{8\pi G_4^{\pm}}{3} T^{\pm}. \quad (3.9)$$

The above equation is very important and relates the curvature on the brane H_{\pm}^2 to its sources, namely the brane tension and the Gauss-Bonnet coupling. We see that the junction conditions tell us that the effective expansion H_{\pm} is in one part due to the Gauss-Bonnet induced cosmological term and in another part due to the vacuum energy of the brane. Since $H_{\pm} = 1/r_{\pm}$, we remind the reader that the Hubble parameter can be expressed as a function of parameters appearing in the soliton potential (2.7). Then, according to the specific explicit solution, the Hubble parameter is constrained and the above relation can give bounds on the two previously mentioned sources of curvature.

A final comment before discussing the various brane world solutions has to do with the induced effective Newton's constant G_4^{\pm} appearing in (3.7). In a usual Kaluza-Klein dimensional reduction, the effective Newton's constant is obtained after substitution of the graviton's zero mode wavefunction in the action and the integration of the extra dimensions. This integration, actually, defines a relation between the effective Newton's constant on the brane, the higher dimensional Newton's constant and the volume of the internal space (see appendix C for the volume calculation for the general brane world models that we will discuss). However, because of the presence of the branes and the Gauss-Bonnet bulk dynamics, the graviton dynamics, as perceived on the brane, have peculiarities and four dimensional dynamics with an induced Newton's constant G_4^{\pm} can be operative, even for internal spaces of infinite volume [8]. This is the interpretation one should give to the effective G_4^{\pm} appearing in (3.7) which can be different from the Newton's constant obtained via the volume calculation in appendix C. For a finite volume element the actual Newton's constant perceived by the brane observer depends on the scale on which we probe gravity on the brane. This is analogous to what happens for the case of the DGP model [8] when we embed it in a Randall-Sundrum setup.

4. Braneworld solutions

In this section, we will analyze some particular cases of brane-world solutions, keeping only some of the parameters in the potential (2.7) each time, which give some interesting brane world examples. In each subcase, we will present the important features introducing the fewest parameters possible. What will interest us in particular are the zeros of V which will correspond to possible brane locations. We will continue to refer to roots of V as horizons although this term is strictly speaking correct only for the black hole solutions (2.6).

4.1 Zero bare cosmological constant

Let us first choose zero bare cosmological constant $a^2 = 0$. Then in all cases of this class, the horizon position is given by the solutions of the algebraic equation

$$x^3 + Ax + M = 0 . \tag{4.1}$$

The roots of the above equation are analyzed in appendix B. In the present section, we will in addition apply the constraints (2.14)–(2.16) and the requirement to be in a spacelike region with $V > 0$, in order to find all the possible vacua.




Figure 1: The horizon positions r_- and r_+ as a function of the black hole mass μ for the Gauss-Bonnet branch $\epsilon = +1$ and for $\alpha < 0$, where the vacua are dS_4 . For $\mu = \mu_c$ we obtain the degenerate horizon case.




Figure 2: The horizon position r_h as a function of the black hole mass μ for the Einstein branch $\epsilon = -1$, where the vacua are dS_4 . The point $\mu = \mu_s$ corresponds to a singularity r_s and is excluded.

Firstly, it is evident that for $a^2 = 0$ there are no $\kappa = 0$ flat vacua since the polynomial (2.16) has no roots². The other cases depend on the branch and the four-dimensional curvature κ .

4.1.1 The dS_4 vacua

Vacua with $\kappa = +1$, which are dS_4 , exist both in the Gauss-Bonnet and the Einstein branches. As we will see, the Gauss-Bonnet branch vacua are compact with respect to the (r, θ) sections, while the Einstein branch vacua are non-compact.

For $\epsilon = +1$, the only solution satisfying all the above requirements is for $\alpha < 0$ and $0 < \mu < \mu_c \equiv 2|\alpha|^{3/2}/(3\sqrt{3})$. Then we have a double root structure, which means that the corresponding brane world solution is of compact (r, θ) sections. It is important to note that in this case the horizon positions are bounded from above $r_- \leq \sqrt{|\alpha|/3}$ and $\sqrt{|\alpha|/3} \leq r_+ \leq \sqrt{|\alpha|}$, so the corresponding Hubble parameters $H_{\pm} = 1/r_{\pm}$ are going to be *bounded from below*. When in particular $\mu = \mu_c$, the two roots become degenerate and $V'_{r_{\pm}} = 0$. The solution deserves special attention and will be discussed later on. The plot of the horizon (or brane) positions as a function of μ is given in figure 1. It is important to stress that the relevant black hole solution corresponding to this soliton,

²This is true for solutions of the specific ansatz (3.2), which does not include the unwarped flat case. As we will see in section 4.1.5, one unwarped flat vacuum for $a^2 = 0$ exists.




Figure 3: The horizon position r_h as a function of the black hole mass μ for the Gauss-Bonnet branch $\epsilon = +1$ and for $\alpha > 0$, where the vacua are AdS_4 .

has the characteristics, of a de-Sitter-Schwarzschild like black hole studied in [31]. Note that the bigger the α the smaller the effective cosmological constant. Here we have a pure Gauss-Bonnet soliton (or black hole) in the sense that the $\alpha \rightarrow 0$ limit is singular.

For $\epsilon = -1$, *i.e.* in the Einstein branch, we have two classes of solutions. The first class is for $\alpha > 0$ and $\mu > 0$. The single horizon of that case can take all positive values. The second class is for $\alpha < 0$ and $\mu > \mu_s \equiv \sqrt{2}|\alpha|^{3/2}$. In this case the horizon position is bounded from below as $r_h > \sqrt{2|\alpha|}$. Furthermore, in the latter case the singularity (2.12) exists and is hidden behind the horizon. In the limit $\mu \rightarrow \mu_s$, we have $r_h \rightarrow r_s$ and the brane position becomes singular. We should therefore exclude the $\mu = \mu_s$ point from the physical parameter space. In both cases, the brane Hubble parameter can be as small as desired. The plots of the horizon position as a function of μ in the above cases is given in figure 2. It is interesting to note that both cases here correspond to a Gauss-Bonnet corrected six dimensional Schwarzschild black hole solution with $\alpha > 0$ or $\alpha < 0$.

4.1.2 The AdS_4 vacua

Vacua with $\kappa = -1$, which are AdS_4 , exist only in the Gauss-Bonnet branch. In the Einstein branch, for all the cases satisfying the constraint (2.15), the region shielded by the horizon is timelike and therefore not suitable for our compactification.

In the Gauss-Bonnet branch $\epsilon = +1$, we have only non-compact brane world models, all of them for $\alpha > 0$. The solutions satisfy the criteria (2.14)–(2.16) for $-\mu_c \leq \mu \leq \mu_b \equiv 5\sqrt{6}\alpha^{3/2}/9$. The singularity (2.12) exists for $0 < \mu \leq \mu_b$ but is always hidden behind the horizon. The plots of the horizon position as a function of μ in the above cases is given in figure 3.

4.1.3 The Nariai dS_4 vacuum

The degenerate case for $\kappa = +1$ in the Gauss-Bonnet branch $\epsilon = +1$ deserves special consideration. This happens when the mass parameter reaches its maximal value $\mu \rightarrow \mu_c$ and so the two horizons become one, $r_- \rightarrow r_+ \rightarrow \sqrt{|\alpha|/3}$. The internal space then appears to collapse, however one should look as for the Nariai metric to a different coordinate system [39]. So, setting $\xi \equiv r_-/r_+$, we make the coordinate transformation that consists

of an affine transformation

$$r = \frac{r_+}{2}[(1 - \xi)R + 1 + \xi], \quad (4.2)$$

$$\theta = \frac{2r_+}{(1 - \xi)}\Theta. \quad (4.3)$$

The degenerate case corresponds to the limit $\xi \rightarrow 1$ and as can be seen from (3.2), it is the limit that the four-dimensional part of the metric is rendered unwarped. In these coordinates $R \in [-1, 1]$ and $\Theta \in [0, 2\pi C)$ with

$$c = \frac{2r_+}{(1 - \xi)}C. \quad (4.4)$$

The affine transformation blows up a point of the coordinate system (r, θ) to a well behaved internal space in the (R, Θ) coordinates. Let us now define the modified potential

$$f = \frac{4V}{(1 - \xi)^2}, \quad (4.5)$$

with the parameters μ and $|\alpha|$ as functions of r_+ and ξ given by

$$\mu = r_+^3 \xi(1 + \xi) \quad , \quad |\alpha| = r_+^2 \frac{1 - \xi^3}{1 - \xi}. \quad (4.6)$$

Then, substituting everything in the metric (3.2) we obtain the blown up metric

$$ds^2 = r_+^2 \left(f d\Theta^2 + \frac{dR^2}{f} + \frac{r^2}{r_+^2} h_{\mu\nu} dx^\mu dx^\nu \right). \quad (4.7)$$

The Nariai limit $\xi \rightarrow 1$ gives a non-singular limit for the modified potential $f \rightarrow \frac{3}{5}(1 - R^2)$. Therefore, we obtain a non-vanishing internal space

$$ds^2 = \left(\frac{|\alpha|}{3} \right) \left[\frac{3}{5}(1 - R^2) d\Theta^2 + \frac{dR^2}{\frac{3}{5}(1 - R^2)} + h_{\mu\nu} dx^\mu dx^\nu \right]. \quad (4.8)$$

Note that, after this coordinate transformation, the deficit angle from (3.5) can be expressed as a function of the modified potential as

$$\beta_+ = \beta_- = \frac{1}{2} |\partial_R f| C = \frac{3}{5} C. \quad (4.9)$$

Thus, in this limit the deficit angle is independent of α and depends only on the periodicity C of the angular coordinate.

4.1.4 Zero black hole mass $\mu = 0$

Let us now suppose that also the black hole mass vanishes $\mu = 0$. Then, the potential has the simple cosmological form

$$V(r) = \kappa + \frac{r^2}{2\alpha}(1 + \epsilon), \quad (4.10)$$

from where we can see that the Einstein branch $\epsilon = -1$ can only give the trivial flat six-dimensional solution, for $\kappa = +1$. The Gauss-Bonnet branch $\epsilon = +1$ on the other hand has non-trivial solutions. The flat $\kappa = 0$ case is excluded since it has no horizon, but there are AdS_4 and dS_4 solutions. Firstly, the $\kappa = -1$ case can have brane world solutions as long as $\alpha > 0$. Then the spacelike region that we use for the internal space is non-compact. This case corresponds to the point of figure 3 where the curve intersects the $\mu = 0$ axis.

On the other hand, for the $\kappa = +1$ vacua we can have an horizon for $\alpha < 0$ and obtain a compact brane world model which is part of a six dimensional de-Sitter space. This case corresponds to the point of figure 1 where the upper curve of r_+ intersects the $\mu = 0$ axis. Furthermore, since the space has no singularity at $r = 0$, we can extend the radial coordinate to $r < 0$ and consider the region of $-\sqrt{|\alpha|} \leq r \leq \sqrt{|\alpha|}$. The internal space is symmetric around $r = 0$, thus we have Z_2 symmetry around the equator of the internal space. The reintroduction of a black hole mass $\mu \neq 0$ breaks this symmetry since it introduces an $r = 0$ singularity.

4.1.5 Flat vacuum

As noted before, the black hole ansatz (3.2) does not admit flat vacua for $a^2 = 0$. However, this ansatz, as we noticed also in the Nariai vacuum, has limitations, since it describes only warped four-dimensional metrics. Looking for unwarped solutions, it is easy to see that the following flat vacuum is a trivial solution of the equations of motion (2.3)

$$ds^2 = \eta_{\mu\nu} dx^\mu dx^\nu + d\rho^2 + \rho^2 d\theta^2, \tag{4.11}$$

where $\theta \in [0, 2\pi c)$. It is non-compact and exists for any sign of α . The equations (3.7), (3.8), (3.9) continue to hold for $\kappa = 0$. To compactify such flat vacua, a gauge field flux has to be added along the lines of appendix A.

4.2 Non-zero bulk cosmological constant

Let us now switch on the bulk cosmological constant, *i.e.* $a^2 \neq 0$. The polynomial (2.16) is more complicated to solve, therefore we will study some special cases and focus on as few parameters as possible. Namely, the case with zero black hole mass $\mu = 0$ and also the point $1 + 4\alpha a^2 = 0$ in parameter space where the theory can be written in a BI form. Finally, we will provide all the flat ($\kappa = 0$) vacua.

4.2.1 Zero black hole mass $\mu = 0$

If the black hole mass vanishes $\mu = 0$, the potential has the simple cosmological form

$$V(r) = \kappa + \frac{r^2}{2\alpha} [1 + \epsilon \sqrt{1 + 4\alpha a^2}]. \tag{4.12}$$

It is obvious that the only brane world solutions that we can obtain in the present case are the curved ones $\kappa = \pm 1$.

Let us first discuss the dS_4 ($\kappa = +1$) solutions. For $\alpha > 0$, we see from (2.15) that only the Einstein branch $\epsilon = -1$ has a solution with a horizon. Furthermore, in this case, in order to have $r_h^2 > 0$, the bulk cosmological constant should be positive, *i.e.* $a^2 > 0$.




Figure 4: The horizon position r_h as a function of the bulk cosmological constant a^2 for the dS_4 vacua with $\mu = 0$. The solution on the left is for $\alpha < 0$, where the Einstein and Gauss Bonnet branches meet at the BI point, represented by dot. The solution on the right is for $\alpha > 0$.




Figure 5: The horizon position r_h as a function of the black hole mass μ for the Born-Infeld case $E = -1$ and for $\alpha > 0$. The left graph is for dS_4 vacua and the right for AdS_4 vacua.

For $\alpha < 0$ we can have solutions both in the Einstein and the Gauss-Bonnet branch. In the Einstein branch $\epsilon = -1$ the horizon is $r_h^2 > 0$ for positive bulk cosmological constant with $0 < a^2 < 1/(4|\alpha|)$. On the other hand, in the Gauss-Bonnet branch $\epsilon = +1$ there are solutions for any $a^2 < 1/(4|\alpha|)$. At the BI limit $a^2 = 1/(4|\alpha|)$ the two branches merge to a unique dS_4 solution. In all the above cases the spacelike region is for $r < r_h$ and the internal spaces are compact. As in section 4.1.4, we can extend the radial coordinate to $r < 0$ and consider the region of $-r_h \leq r \leq r_h$, since the space has no singularity at $r = 0$. The plots of the horizon position as a function of μ in the above cases are given in figure 4.

The AdS_4 ($\kappa = -1$) solutions are the same as above if we substitute $a^2 \rightarrow -a^2$. Thus, the curves in figure 4 should be the mirrors of the $\kappa = 1$ case with respect to the $a^2 = 0$ axis. In addition, the spacelike region for these solutions is for $r > r_h$, thus all the AdS_4 vacua are for non-compact internal spaces.

4.2.2 The Born-Infeld vacua

For the dS_4 and AdS_4 vacua, it is easy to see that V is monotonic for $E = -1$ and has an extremum at $r_{\text{extr}} = (|\alpha|\mu/4)^{1/5}$ if $E = +1$.

The vacua obtained for $1 + 4\alpha a^2 = 0$ deserve special attention. As discussed in the section 2, these vacua do not have an Einstein limit. In a certain way they correspond to the strongly coupled limit of the Gauss-Bonnet term in the action (2.1), since at the linearized level the combination $(1 + 4\alpha a^2)$ multiplies the perturbation operator. Furthermore, as we

saw in the previous section, for these vacua the branches for $\epsilon = +1$, $\epsilon = -1$ merge. Let us now switch on a mass parameter $\mu > 0$ in the potential and write it as

$$V(r) = \kappa + \frac{r^2}{2\alpha} + \frac{M}{\sqrt{r}}, \tag{4.13}$$

where the integration constant $M \equiv \frac{E\sqrt{4|\alpha|\mu}}{2\alpha}$ replaces μ . The potential (4.13) is similar to an Einstein black hole potential in four dimensions with a cosmological constant, apart from the fact that the Newtonian potential is now $1/\sqrt{r}$ rather than $1/r$! Furthermore, if we were to compare it with the six dimensional Einstein black hole ($1/r^3$), we see that gravity in the Born-Infeld case is far weaker as one approaches the singularity.

Keeping this comparison in mind, the flat $\kappa = 0$ vacua of this theory, exist for $E = -1$ and are non-compact. In order for the region $r > r_h$ to be spacelike, we need $\alpha > 0$ and thus $a^2 < 0$ (similar to a planar AdS black hole). The horizon position is simply $r_h = (4\alpha\mu)^{1/5}$. We will see more of the flat vacua in the last subsection.

For the case $E = -1$, the potential has only one root. In order for the region $r > r_h$ to be spacelike, we need to have $\alpha > 0$. Then both cases $\kappa = \pm 1$ are possible. For the AdS_4 ($\kappa = -1$) vacua, the horizon distance is bounded from below $r_h^2 \geq 2\alpha$, while for the dS_4 ($\kappa = +1$) vacua, the horizon is not bounded. The plot of the horizon positions as a function of μ is given in figure 5. The situation is similar to an AdS Schwarzschild black hole with planar or spherical horizon.

In the $E = +1$, the potential has either none or two roots. In order for the potential to acquire two roots the black hole mass should be in the range $\mu > \mu_N \equiv 4(2/5)^{5/2}|\alpha|^{3/2}$. Then we can distinguish two possible cases, one with $\alpha > 0$ and $\kappa = -1$, and a second with $\alpha < 0$ and $\kappa = +1$. For the AdS_4 ($\kappa = -1$) vacua, the spacelike region is for $r > r_+$ and thus the space is non-compact. The horizon is bounded as $2\alpha/5 < r_+^2 < 2\alpha$. For the dS_4 ($\kappa = +1$) vacua, the spacelike region is for $r_- < r < r_+$ and thus the space is compact. The outer horizon is bounded as $2\alpha/5 < r_+^2 < 2\alpha$ and the inner as $0 < r_-^2 < 2\alpha/5$. When $\mu = \mu_N$ the two roots become degenerate and $V'_{r_{\pm}} = 0$. This solution will be discussed in the following section. The plot of the horizon positions as a function of μ is given in figure 6.

4.2.3 The Born-Infeld Nariai vacuum

The degenerate case of the Born-Infeld vacuum for $\kappa = +1$, $E = +1$ and $\alpha < 0$ deserves special consideration. This happens when $\mu \rightarrow \mu_N$ and so $r_- \rightarrow r_+ \rightarrow \sqrt{2|\alpha|/5}$. The internal space then appears to collapse, however one should look as for the Nariai metric in a different coordinate system, as we saw in section 4.1.3. Repeating the same procedure as before and expressing the parameters μ and $|\alpha|$ as functions of r_+ and ξ

$$\mu = \frac{r_+^3}{2} \frac{\xi(1-\xi^2)^2}{(1-\xi^{1/2})(1-\xi^{5/2})}, \quad |\alpha| = \frac{r_+^2}{2} \frac{1-\xi^{5/2}}{1-\xi^{1/2}}, \tag{4.14}$$




Figure 6: The horizon positions r_- and r_+ as a function of the black hole mass μ for the Born-Infeld case $E = +1$ and for $\alpha < 0$, where the vacua are dS_4 . For $\mu = \mu_N$ we obtain the degenerate horizon case. The outer horizon r_+ is also the horizon that we obtain for the case $E = +1$ and for $\alpha > 0$, where the vacua are AdS_4 .

we find that the Nariai limit $\xi \rightarrow 1$ gives a non-singular limit for the modified potential $f \rightarrow \frac{1}{2}(1 - R^2)$. Therefore, we obtain a non-vanishing internal space

$$ds^2 = \left(\frac{2|\alpha|}{5}\right) \left[\frac{1}{2}(1 - R^2)d\Theta^2 + \frac{dR^2}{\frac{1}{2}(1 - R^2)} + h_{\mu\nu}dx^\mu dx^\nu \right]. \quad (4.15)$$

Furthermore, the deficit angle from (3.5) can be expressed as a function of the modified potential as

$$\beta_+ = \beta_- = \frac{1}{2}|\partial_R f|_C = \frac{1}{2}C. \quad (4.16)$$

4.2.4 Flat vacua

Let us now study the flat ($\kappa = 0$) vacua that exist in the $a^2 \neq 0$ case. We have found previously a flat vacuum in the special BI case. More generally, we know from (2.15) that flat vacua can exist only for $\epsilon = -1$. The horizon position is at $r_h^5 = -\mu/a^2$ and we can see from the potential asymptotics that the region $r > r_h$ will be spacelike in two cases: (i) $\alpha < 0$ and $a^2 < 0$ and (ii) $\alpha > 0$ and $-1/(4\alpha) < a^2 < 0$. Thus, for both cases we should have $\mu > 0$. In case (i) there is a branch cut singularity r_s given in (2.12) hidden always beyond the horizon. In case (ii) there is not such r_s .

In all these cases, the flat vacua have non-compact internal spaces and are warped. The latter property is in contrast to the flat vacuum that exists when $a^2 = 0$. Thus, we see that the introduction of a bulk cosmological constant can be compensated by the black hole mass μ and warping of the four-dimensional space, giving again vacua of zero four-dimensional curvature. As noted before, to compactify such flat vacua, a gauge field flux has to be added along the lines of appendix A.

5. Self-properties of the solutions

Let us now discuss the physical consequences of the above solutions. In particular, we wish to see whether we can obtain codimension-2 braneworlds exhibiting *self-accelerating*

or *self-tuning* behaviour. For the former, we seek dS_4 models where the acceleration is mainly of geometrical origin rather than due to the vacuum energy or tension T of the branes. For the latter, we seek braneworld solutions (de Sitter or flat) where the effective Hubble parameter of the brane is not related to the brane vacuum energy T and the tension can be arbitrary large even for small observable Hubble curvature. Here it is important to emphasize, that the Gauss-Bonnet term plays the role of an induced gravity term on the brane [23] according to (3.6). Therefore even if the internal space is not compact we expect that gravity will be four dimensional up to some distance scale. In fact, we would expect, and this turns out to be true, that self-acceleration will hold for warped but uncompactified codimension-2 setups as for the case of codimension-1. Secondly, taking into account that the effective curvature of the brane is given by

$$\kappa H_{\pm}^2 = -\frac{1}{2\alpha} + \frac{8\pi G_4^{\pm}}{3} T^{\pm}, \tag{5.1}$$

i.e. as the sum of a geometrical term and a brane tension term, the self-accelerating and the self-tuning cases are, in some respect, in contrast to each other.

Let us first study *self-acceleration*. The key relation is (5.1). For this case, one should firstly have a non-zero positive geometrical contribution to the curvature, *i.e.* $\alpha < 0$ coming from the Gauss-Bonnet term in the action (2.1). Secondly, this contribution should be dominant in comparison to the brane tension contribution. Note that in order to be in accord with phenomenology from supernovae data [1] and since $H_0 \sim 10^{-34} eV$, the Gauss-Bonnet coupling should be roughly of the order $\alpha \sim 10^{120} M_{Pl}^{-2}$. In this case, the Gauss-Bonnet term in the action is dominant in comparison to the Einstein-Hilbert term. Furthermore, for this limit, even the compact models will have enormous volumes of cosmological size, and therefore will be comparable to certain aspects of infinite volume ones (see appendix C for the calculation of the volumes of the various models).

It is useful to define a self-acceleration index as the fraction of the geometrical acceleration to the curvature

$$s = \frac{1/(2|\alpha|)}{H^2} = \frac{r_h^2}{2|\alpha|}. \tag{5.2}$$

When $0 < s < 1$ we have $T > 0$ assuming $G_4 > 0$. On the other hand, if $s > 1$, we have $T < 0$ with the limit $|T| \rightarrow 3/(16\pi G_4 |\alpha|)$ giving $s \rightarrow \infty$. We can argue that self-acceleration exists whenever $1/2 \lesssim s \lesssim 3/2$, with the strict self-accelerating limit being $s \rightarrow 1$ for $T = 0$.

The index (5.2) depends on the specific bulk solutions that we have. Going through the dS_4 vacua that we discussed in the previous section, we should restrict ourselves to the ones with $\alpha < 0$. Then the self-acceleration index for the various cases is listed as following

1. For $a^2 = 0$, $\epsilon = +1$, $\alpha < 0$, we have two horizons and the index for the two branes varies as $0 < s_- < 1/6$ and $1/6 < s_+ < 1/2$. Thus, for this case (figure 1) we have no strict self-accelerating limit. The Nariai limit case gives $s = 1/6$.
2. For $a^2 = 0$, $\epsilon = -1$, $\alpha < 0$, we have one horizon and the index is $s > 1$ (figure 2). Here we have a strict self-accelerating limit when $\mu \rightarrow \mu_s$, however, at $\mu = \mu_s$ the model is singular.

3. For $a^2 = 0$, $\epsilon = +1$, $\mu = 0$, $\alpha < 0$, we have the compact Z_2 -symmetric model with $s = 1/2$.
4. For $a^2 \neq 0$, $\epsilon = \pm 1$, $\mu = 0$, $\alpha < 0$, we have the compact Z_2 -symmetric model (figure 4). The index for $\epsilon = -1$ is $s > 1$ and for $\epsilon = +1$ varies as $0 < s < 1$. The strict self-accelerating limit happens for the BI case and can be approached by either branch $\epsilon = 1$ and $\epsilon = -1$. One can argue that the BI limit is natural in the sense that the theory becomes more symmetric at that point.
5. For the BI case with $E = +1$, $\mu \neq 0$, $\alpha < 0$, we have two horizons (figure 6) and the index for the two branes varies as $0 < s_- < 1/5$ and $1/5 < s_+ < 1$. The strict self-accelerating limit happens for the $\mu = 0$ case as mentioned above.

From the above we see that the two physically interesting cases where the late acceleration of our Universe can be of purely geometrical origin, are cases 2 and 4. In case 2 the strict $s = 1$ limit can only be reached asymptotically in order not to hit the singularity at $\mu = \mu_s$.

Actually, the first two cases present some similarities and one important difference with the codimension-1 DGP [8], which is worth noting. In fact, we see that case 2 corresponds to the self-accelerating branch of DGP [14] and case 1 to that of the normal branch with dS_4 branes [11]. A first similarity is the fact that we obtain branching, which now is in-between the Gauss-Bonnet branch and the Einstein branch. Secondly, for case 1, as for the normal branch with dS_4 branes, we expect a normalisable spin-2 graviton since the relevant volume element is finite. It is an intriguing difference, however, that the codimension-1 unstable or self-accelerating branch corresponds to the Einstein branch here and not the Gauss-Bonnet one.

In case 4, we note that the limit $s = 1$ is regular and we see that around $a^2 = 1/(2|\alpha|)$, self-acceleration is possible for both branches. Here, the similarity with codimension-1 resides that in that the case of $T = 0$ tension we have enhanced symmetry [10, 11], as one also expects for the Born-Infeld case in codimension-2.

Let us now discuss the cases which have to do with *self-tuning*. The self-tuning idea aims to allow for vacua where their effective cosmological constant is independent from (or at least not strongly dependent on) the vacuum energy of the brane, without fine-tuning. In the toy-model we are presenting here, vacuum energy is represented by the brane tension T_{\pm} whereas the effective cosmological constant is represented by κH_{\pm} . For self-tuning to operate, there should be enough integration constants allowed to be adjusted when we vary one brane tension, with the crucial demand of keeping the curvature of that brane constant. In the solutions that we have discussed, the free parameters are the angular coordinate range c and the black hole mass μ . On the other hand, the action parameters are the Gauss-Bonnet coupling α , the bulk cosmological constant a^2 and the brane tensions T_{\pm} . If we vary for instance T_+ we do not wish α , a^2 , T_- , H_+ to change, but only possibly c and μ . We will be obviously interested in the dS_4 and the flat vacua to test if such a self-tuning can work.

For the dS_4 vacua, the Hubble parameter on the brane depends on the black hole mass μ and the action parameters α and a^2 . Since we require that the action parameters are not tuned, if we fix the curvature H^2 this is equivalent to fixing the black hole mass μ . In the cases where we have more than one brane with different Hubble parameters H_{\pm} , both of the latter are given functions of α , a^2 and μ . Thus, fixing μ from the curvature of the one brane H_+ , fixes also the curvature of the second brane H_- . Then from the relation

$$2\pi(1 - \beta_{\pm}) \left(\frac{1}{2\alpha} + \kappa H_{\pm}^2 \right) = \frac{4\pi G_6}{\alpha} T^{\pm}, \quad (5.3)$$

we see that, if we change T^+ , we can keep the curvature H_+ constant by changing c . But since H_- is fixed, we have to change also T^- . This results to an interbrane fine-tuning. Hence, the only way to obtain self-tuning is when the second brane is absent, or in Z_2 -symmetric models where only one of the above relations survive.

These self-tuning cases, however, are not all theoretically satisfying if we look at the orders of magnitude of the various dimensionful quantities in (5.1). Although we have no idea of the brane tension value today, it would be logical that, if no accurate cancellations happen during phase transitions in the history of the Universe, its natural order of magnitude is $T \sim M_{\text{Pl}}^4$. On the other hand we know that the present value of the Hubble constant is approximately $H_0 \sim 10^{-60} M_{\text{Pl}}$. In order that (5.1) can then be satisfied, today's curvature should be negligible in comparison with $|\alpha|^{-1}$. In other words, we should find models which allow for $|\alpha|H^2 \ll 1$ and approximately

$$\frac{8\pi G_4}{3} T \approx \frac{1}{2\alpha}. \quad (5.4)$$

The above relation should not be viewed as a fine tuning between the action parameters α and T , since G_4 scales as $\frac{1}{\alpha}$ (3.7) and *can vary in time* when the vacuum energy changes, in accordance with the remarks of [35]. The constraints on the time variation of Newton's constant come rather late in the history of the Universe, certainly after the QCD phase transition. Therefore, the relation (5.4) is providing the angle deficit/excess parameter β , given a Gauss-Bonnet coupling α , which needs not to be unnatural as in the self-accelerating case. Furthermore, the relation (5.4) is exact for the case of flat $\kappa = 0$ vacua.

From the models that we have discussed, let us see the cases that give rise to such theoretically viable self-tuning

- For $a^2 = 0$, $\epsilon = -1$, and for both $\alpha > 0$ and $\alpha < 0$, we have the non-compact vacua with $|\alpha|H^2 \ll 1$ for $\mu|\alpha|^{-3/2} \gg 1$ (see figure 2).
- For $a^2 \neq 0$, $\epsilon = -1$, $\mu = 0$, $\alpha < 0$, we have Z_2 -symmetric vacua with $|\alpha|H^2 \ll 1$ for $|\alpha|a^2 \ll 1$ (see figure 4).
- For $a^2 \neq 0$, $\epsilon = -1$, $\mu = 0$, $\alpha > 0$, we have non-compact vacua with $\alpha H^2 \ll 1$ for $\alpha a^2 \ll 1$ (see figure 4).
- For the BI case with $E = -1$, $\alpha > 0$, we have non-compact with $\alpha H^2 \ll 1$ for $\mu\alpha^{-3/2} \gg 1$ (see figure 5).

- All the flat non-compact vacua, for both $\alpha > 0$ and $\alpha < 0$ are self-tuning in the exact sense.

Here we should note once more that the above (exact or approximate) self-tuning solutions do not constitute a resolution, but rather a potential amelioration, of the cosmological constant problem. We have found vacua where the brane curvature is independent of the brane vacuum energy and furthermore the these two energy scales can be well separated. However, the important question, for which the self-tuning mechanism cannot give a simple answer, is if the dynamical variation of the vacuum energy results in remaining in these vacua. In some sense, the term self-tuning itself is deceiving since the existence of these vacua does not guarantee that there are some attractor dynamics which tune the system towards them. Nevertheless, the very existence of them is important, since at least at the level of non-dynamical solutions, dissociating the brane curvature from its vacuum energy is a rather unique situation. Moreover, we saw that most of these vacua are non-compact, but, as we will explain in the last section, gravity is quasi-localised and thus effectively four dimensional up to some cross-over scale.

It is evident that the vacua with the above self-tuning properties are different from the ones with the self-accelerating properties corresponding in particular to a totally different bare parameter α . This is true as long as we want self-acceleration at a very low energy scale. If, for example, we want to explain inflation theory geometrically, as in [36], then the self-accelerating (inflationary) vacua can be deformed to self-tuning ones by letting the integration constant μ run. A nice toy model scenario for this is pictured in figure 2 if we assume that μ is an increasing function of proper cosmological time. One starts at the big-bang singularity which corresponds to a 6 dimensional bulk naked singularity at $\mu = \mu_s$. At early time we have $\mu \gtrsim \mu_s$ and the inflationary expansion is, $H_{\text{inf}}^2 \sim 1/(2|\alpha|)$. At late time the late expansion is related to the large mass of the soliton whereas the vacuum energy is of the order of $1/(G_4|\alpha|)$ which is much larger than the current Universe curvature $H_0^2 M_{\text{Pl}}^2$.

Before we close off this section, it is worth pointing out that no self-tuning cases are possible for the Gauss-Bonnet branch. Furthermore, the only finite volume self-tuning case that we encountered is possible for a de Sitter braneworld.

6. The Gauss-Bonnet instantons

In our previous analysis, we have found several brane world vacua of finite volume. These solutions are of special interest because they can give rise to gravitational instantons. We saw how they all have the feature that the four dimensional space is dS_4 . Wick rotating to a Euclidean metric $h_{\mu\nu}$ as in (2.5) with $\kappa = +1$ and keeping the Euclidean internal space, all the coordinates are spacelike and the solutions describe a Riemannian manifold with conical singularities. Particularly important are the instantons which are regular, in other worlds those having no conical singularities (see also [37, 38]). We can divide these vacua into two classes according to their topology.

The first class of instantons are the ones for vanishing black hole mass $\mu = 0$, where there is no singularity at $r = 0$ and the space is Z_2 -symmetric. These instantons are given

by

$$ds^2 = \left(1 - \frac{r^2}{r_h^2(\alpha, a^2)}\right) d\theta^2 + \frac{dr^2}{\left(1 - \frac{r^2}{r_h^2(\alpha, a^2)}\right)} + r^2 h_{\mu\nu} dx^\mu dx^\nu, \quad (6.1)$$

with $r_h^2 = -2\alpha/(1 + \epsilon\sqrt{1 + 4\alpha a^2})$ and are topologically S^6 . In order for these instantons to be regular, one should have $\beta = 1$, which from (3.5) gives $c = r_h$ for the periodicity of the θ -coordinate. These instantons have an Einstein theory limit only in the case $\epsilon = -1$ and $1 + 4\alpha a^2 \neq 0$.

The second class of instantons are the ones of the Nariai limits of vacua with non-vanishing black hole mass found in section 4.1.3 and section 4.2.3. These instantons are given by

$$ds^2 = r_h^2(\alpha, a^2) \left[\zeta(1 - R^2) d\Theta^2 + \frac{dR^2}{\zeta(1 - R^2)} + h_{\mu\nu} dx^\mu dx^\nu \right], \quad (6.2)$$

where ζ is a numerical factor and r_h the degenerate horizon. These instantons, in contrast with the ones noted before, are topologically $S^2 \times S^4$. In order for these to be regular, one should have $\beta = 1$. The deficit angle for the instantons that we have found is given by $\beta = \zeta C$, thus, regularity of the instantons imposes $C = 1/\zeta$. Let us also note here that these $S^2 \times S^4$ instantons for the case $a^2 \neq 0$, exist also for the case other than the BI limit that we have discussed in section 4.2.3. This is why in (6.2) we allowed for a^2 dependence of the degenerate horizon. None of these instantons has an Einstein theory limit.

The above instantons have the physical interpretation of describing probabilities of nucleation processes [39, 40] between two distinct gravitational backgrounds. In particular, the probability of nucleation of a pair of Nariai black holes from the pure dS_6 backgrounds of $\mu = 0$, is given by

$$\Gamma = \eta \exp[2\Delta S] \quad , \quad \Delta S \equiv S_{S^6} - S_{S^2 \times S^4}, \quad (6.3)$$

where η is the one loop contribution from the quantum quadratic fields and S_{S^6} and $S_{S^2 \times S^4}$ are the values of the action for the two instantons that we presented earlier. It is straightforward to compute the above probability of nucleation for the two simple cases of $a^2 = 0$ and the BI case $a^2 = 1/(4|\alpha|)$. In both cases one obtains

$$\Delta S \propto \frac{\text{Vol}(S^4)}{16\pi G_6} \alpha^2 > 0. \quad (6.4)$$

Since these action differences are positive definite, we deduce that the probability of both processes is unsuppressed unlike the case of [39]. Therefore, the solitonic vacua ($\mu \neq 0$) are apparently stable with respect to pure de Sitter vacua in Lovelock theory. This result certainly deserves further study.

7. Discussion and conclusions

In this paper we studied in some detail codimension-2 braneworlds in the context of Lovelock gravity. Using a modified version of Birkhoff's staticity theorem [24, 25], we found all six dimensional solutions describing a de Sitter, flat or anti de Sitter braneworld of

codimension 2. First thing we can observe is that, unlike the codimension-1 case where the Gauss-Bonnet invariant plays a somewhat secondary role, here, in the case of codimension-2 braneworlds, it can give rise to self-acceleration and certain self-tuning properties which are not present in the Einstein theory. This is largely due to the fact that codimension-2 junction conditions induce on the brane an Einstein-Hilbert term [22]. It is important to note that this does not mean that we have ordinary four-dimensional gravity, rather, as noted in [23], we are in a quite similar situation as for the codimension-1 DGP model [8]. By this, we mean that from some ultraviolet scale up to some infrared scale we expect gravity to "look" four-dimensional. A proof of this statement of course requires the full spectrum of gravitational perturbations for the solutions that we have found.

The important relation we came across is (3.9) which relates the vacuum energy or tension T of the brane with the Gauss-Bonnet coupling α and the effective cosmological expansion on the brane H_0 , which itself is related to the characteristics of the bulk solution: its mass, bulk cosmological constant (and charge) in particular. The interesting feature we found here is that the Gauss-Bonnet coupling can give de Sitter branes without vacuum energy on the brane, purely geometrically. In this sense such de Sitter solutions are self-accelerating. In order to explain the small cosmological constant we then have to fine-tune as usual $\alpha \sim H_0^{-2}$.

The second important point is that Gauss-Bonnet coupling α and the topological parameter β , which is otherwise unconstrained, dictate the cross-over scale between the four-dimensional Planck scale G_4 and the six-dimensional one G_6 (3.7). Here, unlike the five dimensional DGP model, the cross over scale is not necessarily tied up to the self-acceleration scale. Indeed, the more β is close to 1 (3.7), the more we can dissociate these scales. In a nutshell, all depends on the gravity phenomenology beyond the cross-over scale and is dictated by the full graviton propagator on the brane. As we emphasized, the appearance of the induced Einstein tensor in the codimension-2 junction conditions is not a proof of an effective four-dimensional gravity or not. It is a definite sign, however, that a relevant scale will appear in the boundary conditions for the gravitational perturbations. Although we have not undertaken the precise perturbation calculation here, we will comment on some of its characteristics later on.

The most conservative approach is, as in DGP [14], to introduce a hierarchy of scales between G_6 and G_4 so that the cross-over scale is

$$r_c^2 = \frac{G_6}{G_4}. \tag{7.1}$$

Assuming that $r_c \sim H_0^{-1}$, this gives a very low higher dimensional Planck scale $M_6 \equiv G_6^{-1/4} \sim 10^{-30} M_{\text{Pl}}$ which in turn dictates that $1 - \beta \sim \mathcal{O}(1)$. But, as mentioned above, one can have $r_c \ll H_0^{-1}$ by having $1 - \beta \approx 0$. For the self-tuning vacua, on the other hand, the natural value of the Gauss-Bonnet coupling is $\alpha \sim M_{\text{Pl}}^{-2}$ which for the same cross-over scale (7.1) and $r_c \sim H_0^{-1}$ gives a huge angle excess $1 - \beta \sim 10^{120}$. Obviously, in this case a reasonable hierarchy between H_0^{-1} and the cross-over scale r_c , cannot reduce significantly the latter huge excess angle. We emphasize, however, that the above orders of magnitude, should be viewed with caution since the definition of the cross-over scale (7.1)

should be done in a true cosmological setting with complete knowledge of the modified FRW equations.

Another important point concerns the fact that the self-accelerating branch of DGP seems to be embedded in the usual Einstein branch of Lovelock theory and not in the Gauss-Bonnet branch. Indeed, given the Gauss-Bonnet term in the action and no bare cosmological constant ($a^2 = 0$) we find two distinct solutions for $\epsilon = \pm 1$ (2.7): in the Gauss-Bonnet branch, $\epsilon = +1$, we can have a finite volume soliton solution since then the Gauss-Bonnet coupling plays the role of a cosmological constant (figure 1) (situation akin to a Schwarzschild de Sitter black hole [31]). Then, we have a finite volume element and we expect a normalisable zero mode graviton. We get no self-acceleration. For the Einstein branch however, $\epsilon = -1$, we have a single brane, infinite volume element and self acceleration with as small tension as we want (figure 2). The bulk solution corresponds to a Gauss-Bonnet corrected Schwarzschild black hole. This difference may be interesting in respect to linear (in)stability and the presence of ghosts of such backgrounds, not only in the scalar, but also in the spin-2 sector.

Furthermore, in our solutions we see at least two different length scales emerging where we can probe differing four-dimensional gravity. That of the volume element, which if finite, would mean that beyond that scale we expect ordinary four-dimensional gravity. Secondly, that of the cross-over scale where up to that distance gravity seems four-dimensional, as in the case of the codimension-1 DGP model. Again we emphasize that the cross over scale and self acceleration scale are here supplemented by an extra topological parameter β which in a sense tells us how far we are from a Kaluza-Klein setup for the Killing direction ∂_θ . Indeed, we see that for $\beta \rightarrow 0$ our codimension-2 space is very much like a lightning rod [41], where we seemingly have a cascade from a six-dimensional to a five-dimensional and then a four-dimensional setup (reminiscent of the recent proposal [42]).

What is missing from our analysis is the linear perturbation of these solutions which will tell us of the stability and the precise gravitational spectrum. This is not a trivial task, for our metric is not conformally flat. Therefore, apart from the usual complications of black hole perturbations, one has to add the fact that Lovelock perturbations are going to be genuinely different from the usual ones for Einstein theory. The reason is simple: the background Weyl tensor appears in the Lovelock field equations and thus, extra tensor pieces are bound to appear for tensorial perturbations. Recent work in fact from string theory [43], in relation to the AdS/CFT correspondence, indicates that there is yet another scale appearing in the bulk perturbations of (2.6). This extra scale comes from the extra tensorial pieces "to be added" to the usual perturbation operator, that dictate that gravity waves in a planar ($\kappa = 0$) black hole background (2.6) can evolve in a differing background geometry than that of (2.6). This means in particular that the light-cone of the wave fronts can break causality. We think that this is an intriguing and extremely important issue, which we hope will be undertaken soon.

In addition, the cosmology of codimension-2 branes is very poorly understood, especially from the point of view of self-tuning and the issue of vacuum selection. It has been known that in Einstein gravity the introduction of matter other than tension on

codimension-2 branes introduces singularities³ worse than conical [45] (see, however, the work of [46] for intersecting brane cosmology in six dimensions). Although, the appearance of singularities is natural in defects of codimension higher than one [47, 48], one has in practice to regularise the brane [49, 50] in order to consider some cosmological fluid on them [51, 52]. Although one expects that brane singularities persist in the Lovelock theory, the fact that there exists an induced Einstein equation on the brane, allows for the cosmology to be studied without the need of explicit regularisation. We plan to address the question of cosmology of the present Lovelock models in a different publication [53].

Acknowledgments

It is a great pleasure to acknowledge discussions and numerous insightful comments from Nemanja D. Kaloper, Kozuya Koyama and Antonio H. Padilla. We thank them also for reading through the manuscript. We also thank Roy Maartens for bringing up the issue of self acceleration in the context of codimension-2 braneworlds early on. C.C. also thanks Emilian Dudas and Valery Rubakov for discussions and critical comments. A.P. is supported by a Marie Curie Intra-European Fellowship EIF-039189.

A. A more general action

In this appendix we will consider a generalization of the model that we discussed in the main text to a D -dimensional bulk spacetime, where also a gauge field F_{MN} is coupled to gravity. The action of the system reads

$$S = \int d^D x \sqrt{-g} \left[\frac{1}{16\pi G_D} (R + \hat{\alpha} \mathcal{L}_{\text{GB}}) - \Lambda \right] - \frac{1}{4} \int d^D x \sqrt{-g} F^2, \quad (\text{A.1})$$

where G_D the D -dimensional Newton's constant.

It is straightforward to write down the Einstein equations of motion for the above action. They read

$$G_{MN} - \hat{\alpha} H_{MN} = 8\pi G_D T_{MN}, \quad (\text{A.2})$$

with the energy momentum tensor

$$T_{MN} = -\Lambda g_{MN} + F_{MK} F_N^K - \frac{1}{4} F^2 g_{MN}. \quad (\text{A.3})$$

Furthermore, the gauge field equation reads

$$\partial_M (\sqrt{-g} F^{MN}) = 0. \quad (\text{A.4})$$

The static spherically symmetric solutions of the above equations of motion with $(D - 2)$ -dimensional space of maximal symmetry are locally isometric to

$$ds^2 = -V(r) dt^2 + \frac{dr^2}{V(r)} + r^2 h_{\mu\nu} dx^\mu dx^\nu. \quad (\text{A.5})$$

³This is a well known fact in four dimensional gravity, called the Israel paradox concerning self-gravitating cosmic string metrics (see for example [44]).

The explicit solution of the equations of motion then gives

$$F = \frac{q}{4\pi r^{D-2}} dt \wedge dr, \quad (\text{A.6})$$

and for the potential,

$$V(r) = \kappa + \frac{r^2}{2\alpha} \left[1 + \epsilon \sqrt{1 + 4\alpha \left(a^2 - \frac{\epsilon\mu}{r^{D-1}} - \frac{Q^2}{r^{2(D-2)}} \right)} \right], \quad (\text{A.7})$$

and the parameters appearing in the action have been rescaled to,

$$\alpha = (D-3)(D-4)\hat{\alpha}, \quad 16\pi G_D \Lambda = (D-1)(D-2)a^2 \text{ (for } dS_D \text{)}. \quad (\text{A.8})$$

The integration constants are,

$$Q^2 = \frac{G_D q^2}{2\pi(D-2)(D-3)\alpha}, \quad \mu = \frac{16\pi G_D M}{(D-2)\Sigma_\kappa}, \quad (\text{A.9})$$

where q is the charge, M is the AD or ADM mass of the solution and Σ_κ is the area of the unit $(D-2)$ maximally symmetric subspace. Finally, $\epsilon = \pm 1$ which gives rise to 2 branches of solutions. The convention of the μ sign is chosen so that the gravitational mass is always $\mu > 0$. As one can easily check by expanding the square root for large distances, the sign flip in front of μ is necessary to match the Schwarzschild solution behaviour for positive AD mass. On the other hand, the charge term of the above potential is the opposite of a charged black hole for $\epsilon = -1$.

B. The general solution of $x^3 + Ax + M = 0$

To determine the positions of the horizons for the $a^2 = 0$ case, we need the general solutions of the third order algebraic equation

$$x^3 + Ax + M = 0, \quad (\text{B.1})$$

The solutions of the above equation depend on the parameters A , M and in particular on the discriminant of the system

$$D = \left(\frac{M}{2}\right)^2 + \left(\frac{A}{3}\right)^3, \quad (\text{B.2})$$

If $D > 0$ there is only one real root. If $D \leq 0$ there are three real roots, two of which become degenerate where the inequality saturates. A critical value of M that will be useful in the subsequent study is

$$M_c = \frac{2}{3\sqrt{3}}|A|^3. \quad (\text{B.3})$$

Since the solutions of the above equation will correspond to horizons, we also need to look which of these real roots are in addition *positive*. All permissible cases are listed as following




Figure 7: The positive real roots of the equation $x^3 + Ax + M = 0$. The lower curve gives the root for the $A > 0$ case and the upper one the root(s) for the $A < 0$ case. In the interval $0 < M < M_c$, there are two roots $r_- < r_+$ which become degenerate for $M = M_c$.

- If $A > 0$ and $M < 0$, it is $D > 0$ and we have the root

$$r_h = \left[-\frac{M}{2} + \sqrt{D} \right]^{1/3} - \frac{A}{3} \left[-\frac{M}{2} + \sqrt{D} \right]^{-1/3}, \quad (\text{B.4})$$

- If $A < 0$ and $M < -M_c$, it is $D > 0$, but we have one positive root which is given by the same formula as above (B.4). [It has different value, however, since $A < 0$.]
- If $A < 0$ and $0 < M \leq M_c$, it is $D \leq 0$ and we have two roots

$$r_+ = 2\sqrt{\frac{|A|}{3}} \cos \left(\frac{1}{3} \cos^{-1} \left[-\frac{M}{M_c} \right] \right), \quad (\text{B.5})$$

$$r_- = \sqrt{\frac{|A|}{3}} \left\{ -\cos \left(\frac{1}{3} \cos^{-1} \left[-\frac{M}{M_c} \right] \right) + \sqrt{3} \sin \left(\frac{1}{3} \cos^{-1} \left[-\frac{M}{M_c} \right] \right) \right\}. \quad (\text{B.6})$$

- If $A < 0$ and $-M_c \leq M < 0$, it is $D \leq 0$ but we only have one positive root, given by (B.5).

The results of the above analysis are summarized in figure 7.

C. Volume calculation

In this appendix, we will calculate the volume of the general brane world models that we considered in the main text. To do this, we will make the assumption that the zero mode graviton wavefunction follows the warp factor. In other words, we will assume that the zero mode is emanating from the metric

$$ds^2 = V(r)d\theta^2 + \frac{dr^2}{V(r)} + r^2 g_{\mu\nu}^{(4)}(x) dx^\mu dx^\nu, \quad (\text{C.1})$$

with the four dimensional metric $g_{\mu\nu}^{(4)}(x) = h_{\mu\nu} + H_{\mu\nu}$, where $H_{\mu\nu}$ is a perturbation around the background $h_{\mu\nu}$. To find the volume of the space we should substitute the metric (C.1) in the action (2.1) and integrate out the internal space. We will define the volume (Vol) to be given by the coefficient of the four-dimensional curvature term in the action as⁴

$$S = \frac{(\text{Vol})}{16\pi G_6} \int d^4x \sqrt{-g_4} R_4 + \dots, \quad (\text{C.2})$$

where the dots represent higher order in curvature terms. It is straightforward to compute that the various quantities of the action (2.1) for the ansatz (C.1) give

$$\sqrt{-g_6} = r^4 \sqrt{-g_4}, \quad (\text{C.3})$$

$$R_6 = \frac{1}{r^2} R_4 + \dots, \quad (\text{C.4})$$

$$\mathcal{L}_{\text{GB}} = \frac{4}{r^2} \left(\frac{V}{r} \right)' R_4 + \dots \quad (\text{C.5})$$

Then the volume is given by the following integral

$$(\text{Vol}) = 2\pi c \int dr r^2 \left[1 + 4\hat{\alpha} \left(\frac{V}{r} \right)' \right]. \quad (\text{C.6})$$

If $1 + 4\alpha a^2 \neq 0$, the above integral gives

$$(\text{Vol}) = \frac{2\pi c}{3} \left[r^3 - \frac{2\alpha}{3} \left(4\kappa r + \frac{5\mu}{\sqrt{1 + 4\alpha a^2}} \frac{F}{r^2} \right) \right]_{r_-}^{r_+}, \quad (\text{C.7})$$

where F is the hypergeometric function

$$F = {}_2F_1 \left(\frac{2}{5}, \frac{1}{2}, \frac{7}{5}; \frac{4\alpha\epsilon\mu}{(1 + 4\alpha a^2)r^5} \right). \quad (\text{C.8})$$

For the special BI case, $1 + 4\alpha a^2 = 0$, the integral gives instead

$$(\text{Vol}) = \frac{2\pi c}{3} \left[\frac{13}{3} r^3 + 4\alpha\kappa r \right]_{r_-}^{r_+}. \quad (\text{C.9})$$

It is straightforward to see that the above formulas give finite (and positive) volume for the two Nariai cases. To obtain the volume in this case one should move to the (R, Θ) coordinates and take the limit $\xi \rightarrow 1$.

References

- [1] SUPERNOVA SEARCH TEAM collaboration, A.G. Riess et al., *Observational evidence from supernovae for an accelerating universe and a cosmological constant*, *Astron. J.* **116** (1998) 1009 [[astro-ph/9805201](#)];

⁴Notice that according to the ansatz (C.1), it is $[r] = [\theta] = M^{-1}$ and thus $[x^\mu] = M^0$. For this reason, we obtain an unusual mass dimension for the volume $[(\text{Vol})] = M^{-4}$. With a suitable four-dimensional coordinate rescaling, *e.g.* $x_N^\mu = r_h x^\mu$, the volume will be redefined as $(\text{Vol})_N = (\text{Vol})/r_h^2$ and will acquire the correct M^{-2} mass dimension.

- SUPERNOVA COSMOLOGY PROJECT collaboration, S. Perlmutter et al., *Measurements of Omega and Lambda from 42 high-redshift supernovae*, *Astrophys. J.* **517** (1999) 565 [astro-ph/9812133];
- SUPERNOVA SEARCH TEAM collaboration, A.G. Riess et al., *Type Ia supernova discoveries at $z > 1$ from the Hubble space telescope: evidence for past deceleration and constraints on dark energy evolution*, *Astrophys. J.* **607** (2004) 665 [astro-ph/0402512].
- [2] S.M. Carroll, *The cosmological constant*, *Living Rev. Rel.* **4** (2001) 1 [astro-ph/0004075];
 J. Polchinski, *The cosmological constant and the string landscape*, hep-th/0603249;
 R. Bousso, *TASI lectures on the cosmological constant*, *Gen. Rel. Grav.* **40** (2008) 607 [arXiv:0708.4231].
- [3] S. Weinberg, *The cosmological constant problem*, *Rev. Mod. Phys.* **61** (1989) 1.
- [4] S.M. Carroll, *Quintessence and the rest of the world*, *Phys. Rev. Lett.* **81** (1998) 3067 [astro-ph/9806099];
 C.F. Kolda and D.H. Lyth, *Quintessential difficulties*, *Phys. Lett. B* **458** (1999) 197 [hep-ph/9811375];
 N. Kaloper and L. Sorbo, *Of PNGB QuiNtessence*, *JCAP* **04** (2006) 007 [astro-ph/0511543].
- [5] C.M. Will, *The confrontation between general relativity and experiment*, *Living Rev. Rel.* **9** (2005) 3 [gr-qc/0510072]; *Theory and experiment in gravitational physics*, Cambridge University Press, Cambridge U.K. (1993).
- [6] L. Amendola, D. Polarski and S. Tsujikawa, *Power-laws $f(R)$ theories are cosmologically unacceptable*, *Int. J. Mod. Phys. D* **16** (2007) 1555 [astro-ph/0605384].
- [7] I.I. Kogan, S. Mouslopoulos, A. Papazoglou, G.G. Ross and J. Santiago, *A three three-brane universe: new phenomenology for the new millennium?*, *Nucl. Phys. B* **584** (2000) 313 [hep-ph/9912552];
 R. Gregory, V.A. Rubakov and S.M. Sibiryakov, *Opening up extra dimensions at ultra-large scales*, *Phys. Rev. Lett.* **84** (2000) 5928 [hep-th/0002072].
- [8] G.R. Dvali, G. Gabadadze and M. Porrati, *4D gravity on a brane in 5D Minkowski space*, *Phys. Lett. B* **485** (2000) 208 [hep-th/0005016].
- [9] I.I. Kogan, S. Mouslopoulos, A. Papazoglou and L. Pilo, *Radion in multibrane world*, *Nucl. Phys. B* **625** (2002) 179 [hep-th/0105255];
 R. Gregory, V.A. Rubakov and S.M. Sibiryakov, *Gravity and antigravity in a brane world with metastable gravitons*, *Phys. Lett. B* **489** (2000) 203 [hep-th/0003045].
- [10] D. Gorbunov, K. Koyama and S. Sibiryakov, *More on ghosts in DGP model*, *Phys. Rev. D* **73** (2006) 044016 [hep-th/0512097];
 K. Koyama, *Are there ghosts in the self-accelerating brane universe?*, *Phys. Rev. D* **72** (2005) 123511 [hep-th/0503191].
- [11] C. Charmousis, R. Gregory, N. Kaloper and A. Padilla, *DGP spectroscopy*, *JHEP* **10** (2006) 066 [hep-th/0604086].
- [12] G. Dvali, *Predictive power of strong coupling in theories with large distance modified gravity*, *New J. Phys.* **8** (2006) 326 [hep-th/0610013].
- [13] C. Charmousis, R. Gregory and A. Padilla, *Stealth acceleration and modified gravity*, *JCAP* **10** (2007) 006 [arXiv:0706.0857].

- [14] C. Deffayet, *Cosmology on a brane in Minkowski bulk*, *Phys. Lett. B* **502** (2001) 199 [[hep-th/0010186](#)].
- [15] N. Kaloper, *Brane-induced gravity's shocks*, *Phys. Rev. Lett.* **94** (2005) 181601 [*Erratum ibid.* **95** (2005) 059901] [[hep-th/0501028](#)]; *Gravitational shock waves and their scattering in brane-induced gravity*, *Phys. Rev. D* **71** (2005) 086003 [*Erratum ibid.* **71** (2005) 129905] [[hep-th/0502035](#)];
K. Koyama and F.P. Silva, *Non-linear interactions in a cosmological background in the DGP braneworld*, *Phys. Rev. D* **75** (2007) 084040 [[hep-th/0702169](#)].
- [16] C. de Rham and A.J. Tolley, *Mimicking Lambda with a spin-two ghost condensate*, *JCAP* **07** (2006) 004 [[hep-th/0605122](#)].
- [17] N. Arkani-Hamed, S. Dimopoulos, N. Kaloper and R. Sundrum, *A small cosmological constant from a large extra dimension*, *Phys. Lett. B* **480** (2000) 193 [[hep-th/0001197](#)];
S. Kachru, M.B. Schulz and E. Silverstein, *Self-tuning flat domain walls in 5D gravity and string theory*, *Phys. Rev. D* **62** (2000) 045021 [[hep-th/0001206](#)];
S. Förste, Z. Lalak, S. Lavignac and H.P. Nilles, *A comment on self-tuning and vanishing cosmological constant in the brane world*, *Phys. Lett. B* **481** (2000) 360 [[hep-th/0002164](#)];
The cosmological constant problem from a brane-world perspective, *JHEP* **09** (2000) 034 [[hep-th/0006139](#)];
C. Csáki, J. Erlich and C. Grojean, *Gravitational Lorentz violations and adjustment of the cosmological constant in asymmetrically warped spacetimes*, *Nucl. Phys. B* **604** (2001) 312 [[hep-th/0012143](#)];
J.E. Kim, B. Kyae and H.M. Lee, *Self-tuning solution of the cosmological constant problem with antisymmetric tensor field*, *Nucl. Phys. B* **613** (2001) 306 [[hep-th/0101027](#)];
P. Binetruy, C. Charmousis, S.C. Davis and J.-F. Dufaux, *Avoidance of naked singularities in dilatonic brane world scenarios with a Gauss-Bonnet term*, *Phys. Lett. B* **544** (2002) 183 [[hep-th/0206089](#)];
C. Charmousis, S.C. Davis and J.-F. Dufaux, *Scalar brane backgrounds in higher order curvature gravity*, *JHEP* **12** (2003) 029 [[hep-th/0309083](#)].
- [18] J.-W. Chen, M.A. Luty and E. Ponton, *A critical cosmological constant from millimeter extra dimensions*, *JHEP* **09** (2000) 012 [[hep-th/0003067](#)].
- [19] S.M. Carroll and M.M. Guica, *Sidestepping the cosmological constant with football-shaped extra dimensions*, [hep-th/0302067](#);
I. Navarro, *Codimension two compactifications and the cosmological constant problem*, *JCAP* **09** (2003) 004 [[hep-th/0302129](#)];
Y. Aghababaie, C.P. Burgess, S.L. Parameswaran and F. Quevedo, *Towards a naturally small cosmological constant from branes in 6D supergravity*, *Nucl. Phys. B* **680** (2004) 389 [[hep-th/0304256](#)];
I. Navarro, *Spheres, deficit angles and the cosmological constant*, *Class. and Quant. Grav.* **20** (2003) 3603 [[hep-th/0305014](#)];
G.W. Gibbons, R. Güven and C.N. Pope, *3-branes and uniqueness of the Salam-Sezgin vacuum*, *Phys. Lett. B* **595** (2004) 498 [[hep-th/0307238](#)];
J.-M. Schwindt and C. Wetterich, *Holographic branes*, *Phys. Lett. B* **578** (2004) 409 [[hep-th/0309065](#)];
C.P. Burgess, F. Quevedo, G. Tasinato and I. Zavala, *General axisymmetric solutions and self-tuning in 6D chiral gauged supergravity*, *JHEP* **11** (2004) 069 [[hep-th/0408109](#)].

- [20] A. Kehagias, *A conical tear drop as a vacuum-energy drain for the solution of the cosmological constant problem*, *Phys. Lett.* **B 600** (2004) 133 [[hep-th/0406025](#)];
 S. Randjbar-Daemi and V.A. Rubakov, *4D-flat compactifications with brane vorticities*, *JHEP* **10** (2004) 054 [[hep-th/0407176](#)];
 H.M. Lee and A. Papazoglou, *Brane solutions of a spherical σ -model in six dimensions*, *Nucl. Phys.* **B 705** (2005) 152 [[hep-th/0407208](#)];
 A. Kehagias and C. Mattheopoulou, *Flat-brane compactifications in supergravity induced by scalars*, *Nucl. Phys.* **B 797** (2008) 117 [[arXiv:0710.4021](#)].
- [21] D. Lovelock, *The Einstein tensor and its generalizations*, *J. Math. Phys.* **12** (1971) 498;
 B. Zumino, *Gravity theories in more than four-dimensions*, *Phys. Rept.* **137** (1986) 109.
- [22] P. Bostock, R. Gregory, I. Navarro and J. Santiago, *Einstein gravity on the codimension 2 brane?*, *Phys. Rev. Lett.* **92** (2004) 221601 [[hep-th/0311074](#)].
- [23] C. Charmousis and R. Zegers, *Matching conditions for a brane of arbitrary codimension*, *JHEP* **08** (2005) 075 [[hep-th/0502170](#)]; *Einstein gravity on an even codimension brane*, *Phys. Rev.* **D 72** (2005) 064005 [[hep-th/0502171](#)].
- [24] D.L. Wiltshire, *Spherically symmetric solutions of Einstein-Maxwell theory with a Gauss-Bonnet term*, *Phys. Lett.* **B 169** (1986) 36;
 C. Charmousis and J.-F. Dufaux, *General Gauss-Bonnet brane cosmology*, *Class. and Quant. Grav.* **19** (2002) 4671 [[hep-th/0202107](#)].
- [25] R. Zegers, *Birkhoff's theorem in Lovelock gravity*, *J. Math. Phys.* **46** (2005) 072502 [[gr-qc/0505016](#)].
- [26] D.G. Boulware and S. Deser, *String generated gravity models*, *Phys. Rev. Lett.* **55** (1985) 2656.
- [27] R.-G. Cai, *Gauss-Bonnet black holes in AdS spaces*, *Phys. Rev.* **D 65** (2002) 084014 [[hep-th/0109133](#)].
- [28] J. Crisostomo, R. Troncoso and J. Zanelli, *Black hole scan*, *Phys. Rev.* **D 62** (2000) 084013 [[hep-th/0003271](#)].
- [29] S. Deser and B. Tekin, *Energy in generic higher curvature gravity theories*, *Phys. Rev.* **D 67** (2003) 084009 [[hep-th/0212292](#)];
 A. Padilla, *Surface terms and the Gauss-Bonnet Hamiltonian*, *Class. and Quant. Grav.* **20** (2003) 3129 [[gr-qc/0303082](#)].
- [30] R. Gregory and A. Padilla, *Nested braneworlds and strong brane gravity*, *Phys. Rev.* **D 65** (2002) 084013 [[hep-th/0104262](#)].
- [31] R.C. Myers and J.Z. Simon, *Black hole thermodynamics in Lovelock gravity*, *Phys. Rev.* **D 38** (1988) 2434.
- [32] Z.C. Wu, *No-boundary codimension-two braneworld*, *Phys. Lett.* **B 612** (2005) 115 [[hep-th/0405249](#)].
- [33] S. Mukohyama, Y. Sendouda, H. Yoshiguchi and S. Kinoshita, *Warped flux compactification and brane gravity*, *JCAP* **07** (2005) 013 [[hep-th/0506050](#)].
- [34] B. Linet, *The static, cylindrically symmetric strings in general relativity with cosmological constant*, *J. Math. Phys.* **27** (1986) 1817.

- [35] H.-P. Nilles, A. Papazoglou and G. Tasinato, *Selftuning and its footprints*, *Nucl. Phys. B* **677** (2004) 405 [[hep-th/0309042](#)].
- [36] A.A. Starobinsky, *A new type of isotropic cosmological models without singularity*, *Phys. Lett. B* **91** (1980) 99.
- [37] R. Gregory, N. Kaloper, R.C. Myers and A. Padilla, *A new perspective on DGP gravity*, *JHEP* **10** (2007) 069 [[arXiv:0707.2666](#)].
- [38] C. Garraffo, G. Giribet, E. Gravanis and S. Willison, *Gravitational solitons and C^0 vacuum metrics in five-dimensional Lovelock gravity*, [arXiv:0711.2992](#).
- [39] P.H. Ginsparg and M.J. Perry, *Semiclassical perdurance of de Sitter space*, *Nucl. Phys. B* **222** (1983) 245.
- [40] S.W. Hawking, G.T. Horowitz and S.F. Ross, *Entropy, area and black hole pairs*, *Phys. Rev. D* **51** (1995) 4302 [[gr-qc/9409013](#)];
O.J.C. Dias and J.P.S. Lemos, *Pair creation of higher dimensional black holes on a de Sitter background*, *Phys. Rev. D* **70** (2004) 124023 [[hep-th/0410279](#)].
- [41] N. Kaloper and D. Kiley, *Charting the landscape of modified gravity*, *JHEP* **05** (2007) 045 [[hep-th/0703190](#)].
- [42] C. de Rham et al., *Cascading DGP*, [arXiv:0711.2072](#);
C. de Rham, S. Hofmann, J. Khoury and A.J. Tolley, *Cascading gravity and degravitation*, *JCAP* **02** (2008) 011 [[arXiv:0712.2821](#)].
- [43] M. Brigante, H. Liu, R.C. Myers, S. Shenker and S. Yaida, *The viscosity bound and causality violation*, *Phys. Rev. Lett.* **100** (2008) 191601 [[arXiv:0802.3318](#)]; *Viscosity bound violation in higher derivative gravity*, *Phys. Rev. D* **77** (2008) 126006 [[arXiv:0712.0805](#)].
- [44] J.A.G. Vickers, *Generalized cosmic strings*, *Class. and Quant. Grav.* **4** (1987) 1;
V.P. Frolov, W. Israel and W.G. Unruh, *Gravitational fields of straight and circular cosmic strings: relation between gravitational mass, angular deficit, and internal structure*, *Phys. Rev. D* **39** (1989) 1084;
W.G. Unruh, G. Hayward, W. Israel and D. Mcmanus, *Cosmic string loops are straight*, *Phys. Rev. Lett.* **62** (1989) 2897;
B. Boisseau, C. Charmousis and B. Linet, *Dynamics of a self-gravitating thin cosmic string*, *Phys. Rev. D* **55** (1997) 616 [[gr-qc/9607029](#)].
- [45] J.M. Cline, J. Descheneau, M. Giovannini and J. Vinet, *Cosmology of codimension-two braneworlds*, *JHEP* **06** (2003) 048 [[hep-th/0304147](#)].
- [46] H.M. Lee and G. Tasinato, *Cosmology of intersecting brane world models in Gauss-Bonnet gravity*, *JCAP* **04** (2004) 009 [[hep-th/0401221](#)];
O. Corradini, K. Koyama and G. Tasinato, *Induced gravity on intersecting brane-worlds part I: maximally symmetric solutions*, *Phys. Rev. D* **77** (2008) 084006 [[arXiv:0712.0385](#)];
Induced gravity on intersecting brane-worlds part II: cosmology, [arXiv:0803.1850](#).
- [47] C. Charmousis, R. Emparan and R. Gregory, *Self-gravity of brane worlds: a new hierarchy twist*, *JHEP* **05** (2001) 026 [[hep-th/0101198](#)].
- [48] N. Kaloper and D. Kiley, *Exact black holes and gravitational shockwaves on codimension-2 branes*, *JHEP* **03** (2006) 077 [[hep-th/0601110](#)].

- [49] M. Kolanovic, M. Porrati and J.-W. Rombouts, *Regularization of brane induced gravity*, *Phys. Rev. D* **68** (2003) 064018 [[hep-th/0304148](#)];
S. Kanno and J. Soda, *Quasi-thick codimension 2 braneworld*, *JCAP* **07** (2004) 002 [[hep-th/0404207](#)];
I. Navarro and J. Santiago, *Gravity on codimension 2 brane worlds*, *JHEP* **02** (2005) 007 [[hep-th/0411250](#)].
- [50] M. Peloso, L. Sorbo and G. Tasinato, *Standard 4D gravity on a brane in six dimensional flux compactifications*, *Phys. Rev. D* **73** (2006) 104025 [[hep-th/0603026](#)];
E. Papantonopoulos, A. Papazoglou and V. Zamarias, *Regularization of conical singularities in warped six-dimensional compactifications*, *JHEP* **03** (2007) 002 [[hep-th/0611311](#)];
C.P. Burgess, D. Hoover and G. Tasinato, *UV caps and modulus stabilization for 6D gauged chiral supergravity*, *JHEP* **09** (2007) 124 [[arXiv:0705.3212](#)].
- [51] E. Papantonopoulos, A. Papazoglou and V. Zamarias, *Induced cosmology on a regularized brane in six-dimensional flux compactification*, *Nucl. Phys. B* **797** (2008) 520 [[arXiv:0707.1396](#)];
M. Minamitsuji and D. Langlois, *Cosmological evolution of regularized branes in 6D warped flux compactifications*, *Phys. Rev. D* **76** (2007) 084031 [[arXiv:0707.1426](#)].
- [52] J. Vinet and J.M. Cline, *Can codimension-two branes solve the cosmological constant problem?*, *Phys. Rev. D* **70** (2004) 083514 [[hep-th/0406141](#)]; *Codimension-two branes in six-dimensional supergravity and the cosmological constant problem*, *Phys. Rev. D* **71** (2005) 064011 [[hep-th/0501098](#)];
F. Chen, J.M. Cline and S. Kanno, *Modified Friedmann equation and inflation in warped codimension-two braneworld*, *Phys. Rev. D* **77** (2008) 063531 [[arXiv:0801.0226](#)].
- [53] C. Charmousis, G. Kofinas and A. Papazoglou, work in progress.