

Electroweak Symmetry Breaking induced by Dark Matter

Thomas Hambye and Michel H.G. Tytgat¹

¹*Service de Physique Théorique
Université Libre de Bruxelles CP225
Boulevard du Triomphe, 1050 Brussels, Belgium*

The mechanism behind Electroweak Symmetry Breaking (EWSB) and the nature of dark matter (DM) are currently among the most important issues in high energy physics. Since a natural dark matter candidate is a weakly interacting massive particle or WIMP, with mass around the electroweak scale, it is clearly of interest to investigate the possibility that DM and EWSB are closely related. In the context of a very simple extension of the Standard Model, the Inert Doublet Model, we show that dark matter could play a crucial role in the breaking of the electroweak symmetry. In this model, dark matter is the lightest component of an inert scalar doublet. The coupling of the latter with the Standard Model Higgs doublet breaks the electroweak symmetry at one-loop, *à la Coleman-Weinberg*. The abundance of dark matter, the breaking of the electroweak symmetry and the constraints from electroweak precision measurements can all be accommodated by imposing an (exact or approximate) custodial symmetry.

I. INTRODUCTION

One of the goals of the Large Hadron Collider is to elucidate the origin of Electroweak Symmetry Breaking (EWSB). In the framework of the Standard Model (SM), EWSB is expected to be due to the existence of a Brout-Englert-Higgs scalar doublet (Higgs doublet in the sequel) which develops a non-zero vacuum expectation value (v_{ev}) at tree level. This necessitates a negative mass squared for the Higgs doublet, incidentally the only mass term allowed by the symmetries of the SM.

An attractive possibility, proposed long ago by Coleman and Weinberg [1, 2], is that there is no tree level scalar mass altogether –perhaps because of some underlying conformal symmetry– and that EWSB is caused by radiative corrections. However, appealing as it may be, this mechanism fails within the Standard Model. Because of the large negative contribution from top quark loop, either extra gauge bosons [1, 3, 4] or extra scalars [1, 5, 6, 7, 8] with large couplings must be added to the SM to get, within this approach, a Higgs particle mass consistent with the experimental bound [36].

In an apparently different vein, the recent cosmological observations concur to indicate that dark matter exists and that it is even more abundant than ordinary matter [10, 11]. The nature of the dark matter eludes us but a weakly interacting massive particle (WIMP) with mass around the electroweak scale, which was once in thermal equilibrium, would have a relic abundance consistent with observations.

In this article we study a very simple extension of the Standard Model that lies the origin of electroweak symmetry in the existence of a dark matter candidate and its $SU(2)$ partners and their one-loop contribution. This scenario *à la Coleman Weinberg* can give a Higgs mass above the experimental value $M_H > 114.4$ GeV together with a Dark Matter abundance consistent with cosmological observations, $\Omega h^2 \approx 0.12$ [12].

II. THE MODEL

The model we consider is a two Higgs doublet extension of the SM, $H_1 = (h^+ (h + iG_0)/\sqrt{2})^T$ and $H_2 = (H^+ (H_0 +$

$iA_0)/\sqrt{2})^T$, together with a Z_2 symmetry such that all fields of the Standard Model and H_1 are even under Z_2 while $H_2 \rightarrow -H_2$. We assume that Z_2 is not spontaneously broken, *i.e.* that H_2 does not develop a v_{ev} . As there is no mixing between the doublets, h plays the role of the usual Higgs particle. This very minimal extension of the SM is called the Inert Doublet Model (IDM) because the extra (or inert) doublet does not couple to the quarks (and leptons in this version of the model). This feature is consistent with the non-observation of Flavour Changing Neutral Currents.

The IDM has been discussed long ago by Deshpande and Ma [13]. It contains a dark matter candidate in the form of either H_0 or A_0 . This aspect has been considered in recent works, in particular in [14] as a minimal dark matter candidate, in [15] together with a mechanism to generate neutrino masses at one-loop and in [16] as a framework with a heavy Higgs. This dark matter have been further studied in [17, 18, 19]. [37] The most general renormalisable (CP conserving) potential of the model is

$$V = \mu_1^2 |H_1|^2 + \mu_2^2 |H_2|^2 + \lambda_1 |H_1|^4 + \lambda_2 |H_2|^4 + \lambda_3 |H_1|^2 |H_2|^2 + \lambda_4 |H_1^\dagger H_2|^2 + \frac{\lambda_5}{2} [(H_1^\dagger H_2)^2 + h.c.] \quad (1)$$

with real quartic couplings. The $SU(2) \times U(1)$ symmetry is broken by the vacuum expectation value of H_1 , $\langle H_1 \rangle = v/\sqrt{2}$ with $v = -\mu_1^2/\lambda_1 = 246$ GeV while, assuming $\mu_2^2 > 0$, $\langle H_2 \rangle = 0$. The mass of the Higgs boson, h , is

$$m_h^2 = \mu_1^2 + 3\lambda_1 v^2 \equiv -2\mu_1^2 = 2\lambda_1 v^2 \quad (2)$$

while the mass of the charged, H^+ , and two neutral, H_0 and A_0 , components of the field H_2 are given by

$$\begin{aligned} m_{H^+}^2 &= \mu_2^2 + \lambda_3 v^2/2 \\ m_{H_0}^2 &= \mu_2^2 + (\lambda_3 + \lambda_4 + \lambda_5)v^2/2 \\ m_{A_0}^2 &= \mu_2^2 + (\lambda_3 + \lambda_4 - \lambda_5)v^2/2. \end{aligned} \quad (3)$$

Various limits are of interest. There is a Peccei-Quinn symmetry if $\lambda_5 = 0$, with $m_{H_0} = m_{A_0}$. This limit is however disfavoured by constraints from dark matter direct detection experiments [14, 16, 18]. In the limit $\lambda_4 = \lambda_5$, or in the twisted

case $\lambda_4 = -\lambda_5$ [20], there is a custodial $S(3)$ symmetry, with $m_{H^\pm} = m_{A_0}$ or $m_{H^\pm} = m_{H_0}$, respectively. We will come back to the custodial symmetry when we will discuss constraints from LEP precision measurements. Following [16] we parameterise the contribution from symmetry breaking to the mass of H_0 and A_0 by $\lambda_{L,S} = \lambda_3 + \lambda_4 \pm \lambda_5$ (which are also the coupling constants between the Higgs field h and our dark matter candidates H_0 or A_0 respectively).

For appropriate quartic couplings, either H_0 or A_0 is the lightest component of the H_2 doublet and, in absence of other lighter Z_2 -odd fields, either one is a candidate for dark matter. There are *a priori* two distinct dark matter mass scales which have a relic density consistent with WMAP data: a low-mass one, $M_{DM} \lesssim 75$ GeV, below the threshold for W pair production, and a large mass one, $M_{DM} \gtrsim 400$ GeV [14, 16, 18]. The former case is the most promising one from the point of view of direct and/or indirect detection [18, 19]. In this case the DM relic abundance is dictated by a) annihilation of DM into the Higgs, whose efficiency depends on M_h and λ_L or λ_S , b) annihilation into a W^\pm pair, as M_{DM} gets closer to M_{W^\pm} and c) coannihilation of H_0 and A_0 (resp. of DM and H^\pm) into a Z boson (resp. W^\pm), if the mass splitting between H_0 and A_0 (resp. between DM and H^\pm) is, roughly speaking, close to the freeze-out temperature $T_{fo} \sim M_{DM}/20$.

III. ONE-LOOP RADIATIVE CORRECTIONS

We now consider one-loop corrections to the Higgs effective potential, which is given by the usual expression

$$V_{\text{eff}}(h) = \mu_1^2 \frac{h^2}{2} + \lambda_1 \frac{h^4}{4} + \frac{1}{64\pi^2} \sum_i n_i m_i^4 \left(\ln \frac{m_i^2}{\mu^2} - c_i \right) \quad (4)$$

where $n_i = \{1, 1, 1, 1, 2, 2, -12, 2, 4\}$ is the number of degrees of freedom of each species $i = \{h, H_0, G_0, A_0, h^\pm, H^\pm, t, Z, W^\pm\}$ which couples to the Higgs boson with tree level mass (2) and (3) while $m_{G_0}^2 = m_{h^\pm}^2 = \mu_1^2 + \lambda_1 v^2$, $m_t^2 = g_t^2 v^2/2$, $m_W^2 = g^2 v^2/2$ and $m_Z^2 = (g^2 + g'^2)v^2/2$. The constant is $c_i = 3/2$ for all scalars and fermions and $c_i = 5/6$ for all gauge bosons. The gauge bosons loops are given here for completeness. However, as their effects are generically small, we will neglect their contribution in the sequel.

Imposing that the effective potential has an extremum for $\langle h \rangle = v = 246$ GeV, the Higgs mass at one-loop is given by

$$\begin{aligned} M_h^2 &= \frac{d^2 V_{\text{eff}}}{dh^2} = m_h^2 + \frac{1}{32\pi^2} \left[6\lambda_1 f(m_h^2) + \lambda_L f(m_{H_0}^2) \right. \\ &+ 2\lambda_1 f(m_{G_0}^2) + \lambda_S f(m_{A_0}^2) + 4\lambda_1 f(m_{h^\pm}^2) + 2\lambda_3 f(m_{H^\pm}^2) \\ &+ 36\lambda_1^2 h^2 \log \frac{m_h^2}{\mu^2} + \lambda_L^2 h^2 \log \frac{m_{H_0}^2}{\mu^2} + 4\lambda_1^2 h^2 \log \frac{m_{G_0}^2}{\mu^2} \\ &+ \lambda_S^2 h^2 \log \frac{m_{A_0}^2}{\mu^2} + 8\lambda_1^2 h^2 \log \frac{m_{h^\pm}^2}{\mu^2} + 2\lambda_3^2 h^2 \log \frac{m_{H^\pm}^2}{\mu^2} \\ &\left. - 36g_t^2 h^2 f(m_t^2) - 12g_i^4 h^2 \right] \Big|_{\langle h \rangle = v} \quad (5) \end{aligned}$$

with $f(m^2) = m^2(\log(m^2/\mu^2) - 1)$.

Since H_2 has no vacuum expectation value, there is no mixing between the scalars and it is straightforward to compute the contribution of one-loop corrections to the mass of the other scalars from the second derivative of the effective potential around the Higgs vev (see for instance [28], section 11.6). This still requires to keep track of the dependence of the propagators on h , H_0 , A_0 and H^\pm though. The fact that there is no mixing also means that the extremum is necessarily a minimum if all masses are positive. The result is, using the \overline{MS} prescription,

$$\begin{aligned} M_{H_0}^2 &\equiv \frac{\partial^2 V_{\text{eff}}}{\partial H_0^2} = m_{H_0}^2 + \frac{1}{32\pi^2} \left[\lambda_L f(m_h^2) + 6\lambda_2 f(m_{H_0}^2) \right. \\ &+ \lambda_S f(m_{G_0}^2) + 2\lambda_2 f(m_{A_0}^2) + 2\lambda_3 f(m_{h^\pm}^2) + 4\lambda_2 f(m_{H^\pm}^2) \\ &- 2\lambda_L^2 v^2 g(m_h^2, m_{H_0}^2) - 2\lambda_S^2 v^2 g(m_{G_0}^2, m_{A_0}^2) \\ &\left. - (\lambda_4 + \lambda_5)^2 v^2 g(m_{h^\pm}^2, m_{H^\pm}^2) \right] \Big|_{\langle h \rangle = v} \quad (6) \end{aligned}$$

$$\begin{aligned} M_{A_0}^2 &\equiv \frac{\partial^2 V_{\text{eff}}}{\partial A_0^2} = m_{A_0}^2 + \frac{1}{32\pi^2} \left[\lambda_S f(m_h^2) + 2\lambda_2 f(m_{H_0}^2) \right. \\ &+ \lambda_L f(m_{G_0}^2) + 6\lambda_2 f(m_{A_0}^2) + 2\lambda_3 f(m_{h^\pm}^2) + 4\lambda_2 f(m_{H^\pm}^2) \\ &- 2\lambda_S^2 v^2 g(m_h^2, m_{A_0}^2) - 2\lambda_3^2 v^2 g(m_{G_0}^2, m_{H_0}^2) \\ &\left. - (\lambda_4 - \lambda_5)^2 v^2 g(m_{h^\pm}^2, m_{H^\pm}^2) \right] \Big|_{\langle h \rangle = v} \quad (7) \end{aligned}$$

$$\begin{aligned} M_{H^\pm}^2 &\equiv \frac{\partial^2 V_{\text{eff}}}{\partial H^+ \partial H^-} = m_{H^\pm}^2 + \frac{1}{32\pi^2} \left[\lambda_3 f(m_h^2) + 2\lambda_2 f(m_{H_0}^2) \right. \\ &+ \lambda_3 f(m_{G_0}^2) + 2\lambda_2 f(m_{A_0}^2) + 2(\lambda_3 + \lambda_4) f(m_{h^\pm}^2) \\ &+ 8\lambda_2 f(m_{H^\pm}^2) - \frac{1}{2}(\lambda_4 + \lambda_5)^2 v^2 g(m_{h^\pm}^2, m_{H_0}^2) \\ &\left. - 2\lambda_3^2 v^2 g(m_h^2, m_{H^\pm}^2) - \frac{1}{2}(\lambda_4 - \lambda_5)^2 v^2 g(m_{h^\pm}^2, m_{A_0}^2) \right] \Big|_{\langle h \rangle = v} \quad (8) \end{aligned}$$

with $g(m_1^2, m_2^2) = [f(m_1^2) - f(m_2^2)]/(m_2^2 - m_1^2)$.

In all these expressions, we take $\mu = m_t = 172.5$ GeV. In principle, a change in the renormalisation scale is compensated by the scale dependence of the running quartic couplings. However implementing this is a lengthy task since their beta functions mix the different couplings (cf Eq.(61) of appendix B in [16]). At the present exploratory stage we simply neglect the running of the couplings.

IV. EWSB AND DARK MATTER

We first focus on the physically appealing case of vanishing mass terms, or conformal limit $\mu_1 = \mu_2 = 0$. For the sake of completeness, we will comment on the case $\mu_2, \mu_1 \neq 0$ at the end of this section. In the conformal limit there are three important constraints:

1) **EWSB**. The general strategy is simple. The contribution of at least some of the loops with H_2 particles must be large enough to compensate the large, negative, contribution of the top quark. This requires that at least one of the λ_{3-5} couplings must be large and positive. This will inevitably drive

some of the scalar particle masses in the few hundred GeV range. Imagine that EWSB is driven by loop corrections of H^\pm and A_0 , with $\lambda_3 \simeq \lambda_5$. Since these particles represent together only 3 degrees of freedom whereas there are 12 for the top quark, the $\lambda_{3,S}$ contribution is relevant only provided $\lambda_{3,S} \gtrsim 2g_t^2$. Asking that their contribution is large enough for the Higgs mass to be above ~ 115 GeV requires $\lambda_{3,S} \gtrsim 5g_t^2$, approximately. This gives $M_{H^\pm, A_0} \gtrsim 380$ GeV.

2) Low DM mass. In general (see *e.g.* [18]) the mass of DM comes from both μ_2 and the coupling to the Higgs. If $M_{DM} > M_W$, the dominant process for the relic abundance of H_0 is the annihilation into W^\pm and Z pairs. If $\lambda_L = 0$ (or $\lambda_L = 0$), the cross-section scales like $1/M_{DM}^2$ (this is expected on general grounds [30]) and, for a sufficiently large mass, $M_{DM} \gtrsim 400$ GeV, the abundance is consistent with observations. However, if we increase the coupling to the Higgs, it turns out that the DM annihilation cross-section increases and so the DM abundance decreases. (This behaviour is precisely analogous to that of a heavy SM neutrino, whose annihilation cross-section also increases for large neutrino masses [31, 32].) Consequently, if $\mu_2 = 0$ and all the mass comes from the coupling to the Higgs, the abundance of a heavy DM is much smaller than observation and the only viable possibility for dark matter is if $M_{DM} < M_W$.

3) Electroweak precision measurement. Since at least one of the components of the inert doublet must be very heavy to break the electroweak symmetry while the DM candidate must be lighter than M_W , we have to face the constraints from Electroweak Precision Measurements [38]. A doublet with large mass splitting will contribute to the SM ρ parameter or, equivalently, to the Peskin-Takeuchi T parameter. At one-loop

$$\Delta T = \frac{1}{32\pi^2\alpha v^2} [f(M_{H^\pm}, M_{H_0}) + f(M_{H^\pm}, M_{A_0}) - f(M_{A_0}, M_{H_0})] \quad (9)$$

with $f(m_1, m_2) = (m_1^2 + m_2^2)/2 - m_1^2 m_2^2 / (m_1^2 - m_2^2) \ln(m_1^2 / m_2^2)$ [16]. To give an idea, the contribution from $M_{H^\pm} \sim 450$ GeV and $M_{DM} \sim 75$ GeV tree level masses gives $\Delta T \sim 1$ while electroweak precision measurements impose $|\Delta T| \lesssim 0.2$. Since the inert doublet is massless at tree level, strictly speaking ΔT vanishes at one-loop. Nevertheless we should take the issue seriously as the large mass differences we are after will inevitably give a large contribution to the gauge boson mass splitting, be it beyond one-loop order. There is however a nice and painless cure to this problem: as a quick inspection of Eq. (9) reveals, if either H_0 or A_0 is degenerate with H^\pm , the contribution of the inert doublet to the ΔT parameter vanishes identically. Physically, this is due to the existence of a custodial symmetry in the limit $M_{H^\pm} = M_{A_0}$ or $M_{H^\pm} = M_{H_0}$ (*i.e.* $\lambda_4 = \pm\lambda_5$). [39]. Technically, an exact or approximate custodial symmetry does not only avoid large corrections to the T parameter. It also implies that it is no fine tuning to take, for instance, the DM particle to be lighter than the other components of the inert doublet (*i.e.* λ_L or λ_S much different from the other quartic couplings) as required by the EWSB and DM constraints. We think that this feature holds beyond one-loop order.

From the three constraints above, we can now consider four cases (see Table 1). Case I corresponds to a light H_0 and to two

heavy, nearly degenerate A_0 and H^\pm (*i.e.* $m_{H_0} \ll m_{A_0} \simeq m_{H^\pm}$ or $\lambda_L \ll \lambda_S \simeq \lambda_3$). Case II has a reversed hierarchy, *i.e.* $m_{H_0} \lesssim m_{H^\pm} \ll m_{A_0}$ or $\lambda_L \lesssim \lambda_3 \ll \lambda_S$. The two last correspond to A_0 as the DM candidate, with $m_{A_0} \ll m_{H_0} \simeq m_{H^\pm}$ (case III) and $m_{A_0} \lesssim m_{H^\pm} \ll m_{H_0}$ (case IV). Cases III and IV can be obtained from cases I and II simply by switching H_0 with A_0 , *i.e.* λ_5 with $-\lambda_5$. This leaves the relic density unchanged, so that Table 1 is relevant for these cases too.

All the examples of Table 1 have a DM abundance in agreement with WMAP data [40]. As announced, we observe that some of the quartic couplings must be large. Also, in all the working cases the DM mass is below M_W . In Case I (similarly case III), the DM abundance is determined by its annihilation through the Higgs particle only and thus depends on M_h and the effective trilinear hH_0H_0 coupling, *i.e.* $\lambda_L^{eff} = \frac{1}{v} \partial^3 V_{eff} / \partial h \partial^2 H_0 \equiv \frac{1}{v} \partial M_{H_0}^2 / \partial v$ at one-loop. For various, albeit large, couplings we found the correct abundance for DM masses in the range $M_{H_0} \sim (10 - 72)$ GeV. Below this range, the Higgs mediated annihilation is too suppressed. We remark that the values of M_{H_0} consistent with DM and EWSB can be below the ones found in the tree level analysis of [18]. This is because the one-loop contributions to λ_L^{eff} can be sizeable, *i.e.* for the same mass, the DM particle can be more strongly coupled to the Higgs than it is at tree level. In case II (resp. case IV) coannihilation through the W^+ can play a role if the $H^+ - H_0$ (resp. $H^+ - A_0$) splitting is not too large. Notice that the masses of H^\pm quoted in Table 1 are consistent with collider data because the H^+ does not couple to fermions, is short lived and, if $M_{H^\pm} > M_Z/2$, does not contribute to the width of the Z boson. Notice also that, unlike in cases I and III where it plays little role for DM, in cases II and IV the custodial symmetry can only be approximate, otherwise the coannihilation (and direct detection) cross-sections would be too large to be consistent with observations. Notice finally that Cases II and IV require larger quartic coupling because they involve only one heavy degree of freedom in EWSB instead of three in Cases I and III.

Imposing the quartic couplings $\lambda_{3,L,S}$ to be smaller than *e.g.* 2π or 4π gives $M_h \lesssim 80$ GeV or $M_h \lesssim 175$ GeV in Cases II and IV while for Cases I and III we have $M_h \lesssim 150$ GeV or $M_h \lesssim 350$ GeV. We have checked that these M_h bounds can be saturated, keeping $\Omega_{DM} \sim 0.12$. All these numbers are clearly tentative as the quartic couplings are quite large and, even if we are still in the perturbative regime, see *e.g.* Eqs. (16-18) of [16], higher order corrections could be important. However we do not think they would dramatically change the picture drawn here.

It should be clear from these results that, even if $\mu_1, \mu_2 \neq 0$, the existence of DM around the electroweak scale could have startling effects on EWSB. With respect to the conformal limit, larger (smaller) quartic coupling than in the conformal case should be considered for $\mu_2^2 > 0$ (resp. $\mu_2^2 < 0$).

Instead of an inert doublet we could consider higher-dimensional inert Higgs multiplets. The case of a scalar singlet has already been considered to induce EWSB [7, 34]. In our opinion, in the latter case the connection discussed in the present paper would be looser since it could not be the same object that drive both the EWSB and has the right relic DM

	λ_1	λ_2	λ_3	λ_4	λ_5	M_h	M_{H_0}	M_{A_0}	M_{H^\pm}	h_{BR}	W_{BR}
I	-0.11	0	5.4	-2.8	-2.8	120	12	405	405	100%	0%
I	-0.11	-2	5.4	-2.7	-2.7	120	43	395	395	100%	0%
I	-0.11	-3	5.4	-2.6	-2.6	120	72	390	390	94%	6%
I	-0.30	0	7.6	-4.1	-4.1	180	12	495	495	100%	0%
I	-0.30	-2.5	7.6	-3.8	-3.8	180	64	470	470	100%	0%
II	-0.18	-3	-0.003	4.6	-4.7	120	39	500	55	100%	0%
II	-0.29	-5	-0.07	5.5	-5.53	150	54	535	63	0%	100%

TABLE I: Instances of parameters with WMAP DM abundance. Also given are the relative contribution of Higgs mediated annihilation (h_{BR}) and gauge processes (W_{BR}).

abundance. Assuming their masses to be around the EW scale, several singlets with large (for EWSB) and small couplings (for DM [35]) would presumably be necessary.

V. SUMMARY

We have shown that Dark Matter in the form of the lightest neutral component of a single inert doublet could be responsible for EWSB. We have met essentially three constraints. A

large quartic coupling is necessary to drive EWSB. One quartic coupling must be small to have a DM particle mass below M_W . Finally a small mass splitting of either the A_0 or H_0 with H^\pm is required to confront Electroweak Precision Measurements. All these conditions can be satisfied naturally if an exact or approximate custodial symmetry is assumed. As a result of all constraints we get the bound on the mass of the Higgs $M_h \lesssim 350$ GeV while the mass of dark matter is in the range $M_{DM} \sim (10 - 72)$ GeV. Such a DM candidate is in a range of couplings that makes it accessible to both direct (ZEPLIN) and indirect (GLAST) future searches (cf Figure 5 of [18]).

Since the quartic couplings are quite large, the results of the present paper are probably only tentative. Nevertheless we do expect that the breaking of the Electroweak Symmetry with a WIMP Dark Matter candidate is a feature of the Inert Doublet Model which will survive further investigations.

Acknowledgement

We thank J.-M. Gérard for useful discussions. This work is supported by the FNRS, the IISN and the Belgian Federal Science Policy (IAP VI/11). Preprint ULB-TH/07-26.

-
- [1] S. R. Coleman and E. Weinberg, Phys. Rev. D **7** (1973) 1888.
[2] M. Sher, Phys. Rept. **179**, 273 (1989).
[3] R. Hempfling, Phys. Lett. B **379** (1996) 153.
[4] W. F. S. Chang, J. N. Ng and J. M. S. Wu, arXiv:hep-ph/0701254.
[5] E. Gildener and S. Weinberg, Phys. Rev. D **13**, 3333 (1976).
[6] K. A. Meissner and H. Nicolai, Phys. Lett. B **648**, 312 (2007).
[7] J. R. Espinosa and M. Quiros, arXiv:hep-ph/0701145.
[8] R. Foot, A. Kobakhidze and R. R. Volkas, arXiv:0704.1165 [hep-ph].
[9] T. Hambye, Phys. Lett. B **371**, 87 (1996).
[10] D. N. Spergel *et al.*, arXiv:astro-ph/0603449.
[11] U. Seljak, A. Slosar and P. McDonald, JCAP **0610** (2006) 014.
[12] W. M. Yao *et al.* [Particle Data Group], J. Phys. G **33** (2006) 1.
[13] N. G. Deshpande and E. Ma, Phys. Rev. D **18** (1978) 2574.
[14] M. Cirelli, N. Fornengo and A. Strumia, Nucl. Phys. **B753** (2006) 178.
[15] E. Ma, Phys. Rev. D **73** (2006) 077301.
[16] R. Barbieri, L. J. Hall and V. S. Rychkov, Phys. Rev. D **74** (2006) 015007 [arXiv:hep-ph/0603188].
[17] D. Majumdar and A. Ghosal, arXiv:hep-ph/0607067.
[18] L. Lopez Honorez, E. Nezri, J. F. Oliver and M. H. G. Tytgat, JCAP **0702** (2007) 028
[19] M. Gustafsson, E. Lundstrom, L. Bergstrom and J. Edsjo, arXiv:astro-ph/0703512.
[20] J. M. Gérard and M. Herquet, arXiv:hep-ph/0703051.
[21] A. Pierce and J. Thaler, arXiv:hep-ph/0703056.
[22] M. Lisanti and J. G. Wacker, arXiv:0704.2816 [hep-ph].
[23] X. Calmet and J. F. Oliver, Europhys. Lett. **77** (2007) 51002.
[24] J. A. Casas, J. R. Espinosa and I. Hidalgo, arXiv:hep-ph/0607279.
[25] J. Kubo, E. Ma and D. Suematsu, Phys. Lett. **B642** (2006) 18.
[26] T. Hambye, K. Kannike, E. Ma and M. Raidal, Phys. Rev. **D75** (2007) 095003.
[27] E. Ma, Mod. Phys. Lett. A **21** (2006) 1777.
[28] M. E. Peskin and D. V. Schroeder, *Reading, USA: Addison-Wesley (1995) 842 p*
[29] G. Jungman, M. Kamionkowski and K. Griest, Phys. Rept. **267** (1996) 195.
[30] K. Griest and M. Kamionkowski, Phys. Rev. Lett. **64** (1990) 615.
[31] K. Enqvist, K. Kainulainen and J. Maalampi, Nucl. Phys. B **317** (1989) 647.
[32] K. Kainulainen and K. A. Olive, arXiv:hep-ph/0206163.
[33] G. Belanger, F. Boudjema, A. Pukhov and A. Semenov, Comput. Phys. Commun. **176** (2007) 367.
[34] B. Patt and F. Wilczek, arXiv:hep-ph/0605188.
[35] J. McDonald, Phys. Rev. Lett. **88** (2002) 091304.
[36] Another possibility, that we will not address here, is to consider the SM as an effective theory cutted off by a hard scale [9].
[37] Variations on the IDM from various perspectives has been discussed in e.g. [20, 21, 22, 23, 24, 25, 26, 27].
[38] In [16] the mass splitting must be kept small $\lesssim 15$ GeV because the Higgs is assumed to be very heavy $M_h \gtrsim 500$ GeV and the abundance is dictated by coannihilation. This does not apply in our case because annihilation goes through the Higgs.
[39] Notice that the hypothesis of custodial symmetry together with the Z_2 symmetry automatically gives CP conservation in the scalar sector [20]. CP violation (*i.e.* λ_5 complex) is potentially dangerous for dark matter. It leads to H_0 - A_0 mixing and thus to the possibility of spin-independent direct detection through Z-exchange. Unless the phase of λ_5 is tiny, this induces a far too large direct detection rate for WMAP abundances.
[40] The relic abundance was computed using Micromegas2.0 [33]