

Evidence for a Protophobic Fifth Force from ${}^8\text{Be}$ Nuclear Transitions

Jonathan L. Feng,¹ Bartosz Fornal,¹ Iftah Galon,¹ Susan Gardner,^{1,2}
Jordan Smolinsky,¹ Tim M. P. Tait,¹ and Philip Tanedo¹

¹*Department of Physics and Astronomy, University of California, Irvine, California 92697-4575 USA*
²*Department of Physics and Astronomy, University of Kentucky, Lexington, Kentucky 40506-0055 USA*

Recently a 6.8σ anomaly has been reported in the opening angle and invariant mass distributions of e^+e^- pairs produced in ${}^8\text{Be}$ nuclear transitions. The data are explained by a 17 MeV vector gauge boson X that is produced in the decay of an excited state to the ground state, ${}^8\text{Be}^* \rightarrow {}^8\text{Be}X$, and then decays through $X \rightarrow e^+e^-$. The X boson mediates a fifth force with a characteristic range of 12 fm and has milli-charged couplings to up and down quarks and electrons, and a proton coupling that is suppressed relative to neutrons. The protophobic X boson may also alleviate the current 3.6σ discrepancy between the predicted and measured values of the muon's anomalous magnetic moment.

PACS numbers: 14.70.Pw, 27.20.+n, 21.30.-x, 12.60.Cn, 13.60.-r

Introduction. The four known forces of nature, the electromagnetic, weak, strong, and gravitational interactions, are mediated by the photon, the W and Z bosons, the gluon, and the graviton, respectively. The possibility of a fifth force, similarly mediated by an as-yet-unknown gauge boson, has been discussed [1] since shortly after the introduction of Yang-Mills gauge theories, and has a rich, if checkered, history [2]. If such a force exists, it must either be weak, or short-ranged, or both to be consistent with the wealth of experimental data. In recent years, interest in this possibility has been heightened by the obvious need for dark matter, which has motivated new particles and forces in a dark or hidden sector that may mix with the visible sector and naturally induce a weak fifth force between the known particles.

Recently, studies of decays of an excited state of ${}^8\text{Be}$ to its ground state have found a 6.8σ anomaly in the opening angle and invariant mass distribution of e^+e^- pairs produced in these transitions [3]. The discrepancy from expectations may be explained by as-yet-unidentified nuclear reactions or experimental effects, but the observed distribution is beautifully fit by assuming the production of a new boson. In this work, we advance the new particle interpretation, carefully considering the putative signal and the many competing constraints on its properties, and present a viable proposal for the new boson and the fifth force it induces.

The ${}^8\text{Be}$ Decay Anomaly. The ${}^8\text{Be}$ nuclear excitation spectrum is precisely known [4]. For this discussion, the most relevant ${}^8\text{Be}$ nuclear states and their properties are given in Table I. To simplify our notation, we use the given symbols to denote specific states. The ground state atomic mass is $8.005305\text{ u} \simeq 7456.89\text{ MeV}$; the ground state nuclear mass listed in Table I is about $4m_e$ below this. There are also several unlisted broad resonance excited states both above and below ${}^8\text{Be}^*$ and ${}^8\text{Be}^{*'}$ with widths as large as several MeV.

In the experiment of Krasznahorkay *et al.* [3], an intense proton beam impinges on thin ${}^7\text{Li}$ targets. Given

TABLE I. Relevant ${}^8\text{Be}$ states and their masses, decay widths, and spin-parity and isospin quantum numbers.

State	Mass (MeV)	Width (keV)	J^P	Isospin
${}^8\text{Be}^*$ (18.15)	7473.00	138	1^+	0
${}^8\text{Be}^{*'}$ (17.64)	7472.49	10.7	1^+	1
${}^8\text{Be}$ (g.s.)	7454.85	—	0^+	0

the ${}^7\text{Li}$ nucleus mass of 6533.83 MeV, the ${}^8\text{Be}^*$ and ${}^8\text{Be}^{*'}$ states are resonantly produced by tuning the proton kinetic energies to 1.025 and 0.441 MeV, respectively. The resulting excited states then decay promptly, dominantly back to $p{}^7\text{Li}$, but also through rare electromagnetic processes. For ${}^8\text{Be}^*$, radiative decay to the ground state has branching ratio $B({}^8\text{Be}^* \rightarrow {}^8\text{Be}\gamma) \approx 1.4 \times 10^{-5}$, and there are also decays via internal pair conversion (IPC) with branching ratio $B({}^8\text{Be}^* \rightarrow {}^8\text{Be}e^+e^-) \approx 3.9 \times 10^{-3} B({}^8\text{Be}^* \rightarrow {}^8\text{Be}\gamma) \approx 5.5 \times 10^{-8}$ [5].

For the IPC decays, one can measure the opening angle Θ between the e^+ and e^- and also the invariant mass $m_{e^+e^-}$. One expects these distributions to be sharply peaked at low values of Θ and $m_{e^+e^-}$ and fall smoothly and monotonically for increasing values. This is not what is seen in the ${}^8\text{Be}^*$ decays. Instead, there are pronounced bumps at $\Theta \approx 140^\circ$ and at $m_{e^+e^-} \approx 17\text{ MeV}$ [3]. The experimental analysis fits the contributions from nearby broad resonances, but these cannot reproduce the shape of the observed excesses. The deviation has a significance of 6.8σ , corresponding to a background fluctuation probability of 5.6×10^{-12} [3]. The excess is maximal on the ${}^8\text{Be}^*$ resonance and disappears as the proton beam energy is moved off resonance. No such effect is seen in ${}^8\text{Be}^{*'}$ IPC decays.

The fit may be improved by postulating a new boson X that is produced on-shell in ${}^8\text{Be}^* \rightarrow {}^8\text{Be}X$ and decays promptly via $X \rightarrow e^+e^-$. The authors of Ref. [3] have simulated this process, including the detector energy resolution, which broadens the $m_{e^+e^-}$ peak significantly [6]. They find that the observed excess's shape and

size are beautifully fit by a new boson with mass $m_X = 16.7 \pm 0.35$ (stat) ± 0.5 (sys) MeV and relative branching ratio $B(^8\text{Be}^* \rightarrow ^8\text{Be} X)/B(^8\text{Be}^* \rightarrow ^8\text{Be} \gamma) = 5.6 \times 10^{-6}$, assuming $B(X \rightarrow e^+e^-) = 1$. With these values, the fit had a $\chi^2/\text{dof} = 1.07$.

Protophobic Gauge Bosons. *A priori* the X boson may be a scalar, pseudoscalar, vector, axial vector, or even a spin-2 particle. Some of these cases are easy to dismiss. If parity is conserved, the X boson cannot be a scalar: in a $1^+ \rightarrow 0^+0^+$ transition, angular momentum conservation requires the final state to have $L = 1$, but parity conservation requires $+1 = (-1)^L$. Decays to a pseudoscalar 0^- state are not forbidden by any symmetry, but are severely constrained by experiment. For such axion-like particles a , the two-photon interaction $g_{a\gamma\gamma} a F^{\mu\nu} \tilde{F}_{\mu\nu}$ is almost certainly present at some level, but for $m_a \approx 17$ MeV, all coupling values in the range $1/(10^{18} \text{ GeV}) < g_{a\gamma\gamma} < 1/(10 \text{ GeV})$ are excluded [7, 8].

Here we focus on the vector case. We consider a massive spin-1 Abelian gauge boson X that couples non-chirally to standard model (SM) fermions with charges ε_f in units of e . The new Lagrangian terms are

$$\mathcal{L} = -\frac{1}{4} X_{\mu\nu} X^{\mu\nu} + \frac{1}{2} m_X^2 X_\mu X^\mu - X^\mu J_\mu, \quad (1)$$

where X has field strength $X_{\mu\nu}$ and couples to the current $J_\mu = \sum_f e \varepsilon_f \bar{f} \gamma_\mu f$, or, at the nucleon level, $J_\mu^N = e \varepsilon_p \bar{p} \gamma_\mu p + e \varepsilon_n \bar{n} \gamma_\mu n$, with $\varepsilon_p = 2\varepsilon_u + \varepsilon_d$ and $\varepsilon_n = \varepsilon_u + 2\varepsilon_d$.

We first determine what values of the charges are required to fit the ^8Be signal. The characteristic energy scale of the decay $^8\text{Be}^* \rightarrow ^8\text{Be} X$ is 10 MeV, and so we may consider an effective theory in which $^8\text{Be}^*$, ^8Be , and X are the fundamental degrees of freedom. The one effective operator consistent with the J^P quantum numbers of these states is

$$\mathcal{L}_{\text{int}} = \frac{1}{\Lambda} \varepsilon^{\mu\nu\alpha\beta} (\partial_\mu ^8\text{Be}_\nu^* - \partial_\nu ^8\text{Be}_\mu^*) X_{\alpha\beta} ^8\text{Be}. \quad (2)$$

The matrix element $\langle ^8\text{Be} X | \mathcal{L}_{\text{int}} | ^8\text{Be}^* \rangle$ is proportional to $\langle ^8\text{Be} | J_\mu^N | ^8\text{Be}^* \rangle = (e/2)(\varepsilon_p + \varepsilon_n) \mathcal{M}$, where $\mathcal{M} = \langle ^8\text{Be} | (\bar{p} \gamma_\mu p + \bar{n} \gamma_\mu n) | ^8\text{Be}^* \rangle$ contains the isoscalar component of the current, since the initial and final states are both isoscalars. The resulting decay width is

$$\Gamma(^8\text{Be}^* \rightarrow ^8\text{Be} X) = \frac{(e/2)^2 (\varepsilon_p + \varepsilon_n)^2}{3\pi \Lambda^2} |\mathcal{M}|^2 |\vec{p}_X|^3. \quad (3)$$

To fit the signal, we need

$$\frac{B(^8\text{Be}^* \rightarrow ^8\text{Be} X)}{B(^8\text{Be}^* \rightarrow ^8\text{Be} \gamma)} = (\varepsilon_p + \varepsilon_n)^2 \frac{|\vec{p}_X|^3}{|\vec{p}_\gamma|^3} \approx 5.6 \times 10^{-6}, \quad (4)$$

where both the nuclear matrix elements and the scale Λ have canceled in the ratio. For $m_X = 17$ MeV, we require $|\varepsilon_p + \varepsilon_n| \approx 0.011$, or

$$|\varepsilon_u + \varepsilon_d| \approx 3.7 \times 10^{-3}. \quad (5)$$

The 17 MeV X boson is produced through hadronic couplings, but can decay only to e^+e^- , $\nu\bar{\nu}$, or $\gamma\gamma\gamma$. (We assume there are no decays to unknown particles.) The three-photon decay is negligible, and we will assume that decays to neutrinos are also highly suppressed, for reasons given below. The X boson then decays through its electron coupling with width [9]

$$\Gamma(X \rightarrow e^+e^-) = \varepsilon_e^2 \alpha \frac{m_X^2 + 2m_e^2}{3m_X} \sqrt{1 - 4m_e^2/m_X^2}. \quad (6)$$

The X boson is produced with velocity $v \approx 0.35c$ in the $^8\text{Be}^*$ frame, which is moving non-relativistically with $v = 0.017c$ relative to the lab frame. The X mean decay length is $L \approx \varepsilon_e^{-2} 1.8 \times 10^{-12}$ m in the lab frame. The X boson must decay promptly in the experimental setup of Refs. [3, 6] so that the e^+e^- decay products are detected and the Θ measurements are not distorted. Requiring $L \lesssim 1$ cm, for example, implies

$$|\varepsilon_e| \gtrsim 1.3 \times 10^{-5}. \quad (7)$$

From Eq. (5), we see that a dark photon cannot explain the ^8Be anomaly. For a dark photon, fermions have charges proportional to their SM charges, $\varepsilon_f = q_f \varepsilon$, where ε is the kinetic mixing parameter, and so Eq. (5) implies $\varepsilon \approx 0.011$. This is excluded by many experiments, and most stringently by NA48/2, which requires $\varepsilon < \varepsilon_{\text{max}} = 8 \times 10^{-4}$ at 90% CL [10]. The authors of Ref. [3] estimated that $\varepsilon^2 \sim 10^{-7}$ can fit the signal, but this value of ε is far too small, in part because of the $|\vec{p}|^3$ suppression of the signal.

The NA48/2 bound, however, does not exclude a general vector boson interpretation of the ^8Be anomaly. The NA48/2 limit is a bound on $\pi^0 \rightarrow X\gamma$. In the general gauge boson case, this is proportional to the anomaly trace factor $N_\pi \equiv (\varepsilon_u q_u - \varepsilon_d q_d)^2$. Applying the dark photon bound $N_\pi < \varepsilon_{\text{max}}^2/9$, we find that, for a general gauge boson,

$$|2\varepsilon_u + \varepsilon_d| < \varepsilon_{\text{max}} = 8 \times 10^{-4}. \quad (8)$$

Equations (5) and (8) may be satisfied with a mild $\sim 10\%$ cancelation, provided the charges satisfy

$$-2.3 < \frac{\varepsilon_d}{\varepsilon_u} < -1.8, \quad -0.067 < \frac{\varepsilon_p}{\varepsilon_n} < 0.078. \quad (9)$$

Given the latter condition, we call the general class of vector models that can both explain the ^8Be anomaly and satisfy pion decay constraints ‘‘protophobic.’’

Constraints from Other Experiments. Although there is no need for the gauge boson to decouple from protons completely, for simplicity, for the rest of this work, we consider the extreme protophobic limit where $\varepsilon_p = 0$. We parameterize the quark charges as $\varepsilon_u = -\frac{1}{3}\varepsilon_n$, $\varepsilon_d = \frac{2}{3}\varepsilon_n$ and determine what choices for ε_n , ε_e , and ε_ν are viable. We focus on these first-generation charges, as the




FIG. 1. The required charges to explain the ^8Be anomaly in the (ϵ_u, ϵ_d) (top) and $(\epsilon_e, \epsilon_\nu)$ (bottom) planes, along with the leading constraints discussed in the text. *Top*: The n -Pb and NA48/2 constraints are satisfied in the shaded regions. On the protophobic contour, $\epsilon_d/\epsilon_u = -2$. The width of the ^8Be bands corresponds to requiring the signal strength to be within a factor of 2 of the best fit. *Bottom*: The E141, KLOE-2, $(g-2)_e$, and $\nu - e$ scattering constraints exclude their shaded regions, whereas $(g-2)_\mu$ favors its shaded region. The ^8Be signal imposes a lower bound on $|\epsilon_e|$.

^8Be signal depends on them, but include comments on the charges of the other generations below. The charges required to explain the ^8Be signal, along with the leading bounds discussed below, are shown in Fig. 1.

As noted above, the decay $^8\text{Be}^{*'} \rightarrow ^8\text{Be} X$ is not seen. The protophobic gauge boson can mediate isovector transitions, so there is no dynamical suppression of this decay. However, its mass is near the 17.64 MeV threshold, so the decay is kinematically suppressed. For

$m_X = 17.0$ (17.4) MeV, the $|\vec{p}_X|^3/|\vec{p}_\gamma|^3$ phase space suppression factor is 2.3 (5.2) times more severe for the $^8\text{Be}^{*'}$ decay than for the $^8\text{Be}^*$ decay. In particular, $m_X = 17.4$ MeV is within 1σ of the central value, and a 5.2 times smaller signal in the $^8\text{Be}^{*'}$ decay is consistent with the data. We will continue to refer to the boson as a 17 MeV boson, as no other processes are sensitive to the precise value of its mass, with the understanding that the null $^8\text{Be}^{*'}$ result may require it to be a bit above 17 MeV. Note that although $m_X = 17.4$ MeV is near the endpoint of the $^8\text{Be}^{*'}$ decay, it is not near the endpoint of the $^8\text{Be}^*$ decay, and the Θ and $m_{e^+e^-}$ distributions return to near their SM values at high values. This is not a “last bin” effect.

A number of experiments provide upper bounds on $|\epsilon_e|$. The anomalous magnetic moment of the electron, $(g-2)_e$, constrains $|\epsilon_e| < 1.4 \times 10^{-3}$ (3σ) [11]. The KLOE-2 experiment has looked for $e^+e^- \rightarrow \gamma X$, followed by $X \rightarrow e^+e^-$, and finds $|\epsilon_e| < 2 \times 10^{-3}$ [12]. A similar search at BaBar has reached similar sensitivity in ϵ_e , but is limited to $m_X > 20$ MeV [13].

Electron beam dump experiments also constrain ϵ_e by searching for X bosons radiated off electrons that scatter on target nuclei. As a group, these exclude $|\epsilon_e|$ in the 10^{-8} to 10^{-4} range [14]. For this discussion, given Eq. (7), these experiments provide lower bounds on $|\epsilon_e|$. In more detail, for $m_X \approx 17$ MeV, SLAC experiment E141 requires $|\epsilon_e| > 2 \times 10^{-4}$ [15, 16]. There are also less stringent bounds from Orsay [17] and SLAC’s E137 [18] and Millicharge [19] experiments, and Fermilab experiment E774 [20] excludes some couplings when $m_X < 10$ MeV.

We now turn to bounds on the hadronic couplings. We have already discussed the bound of NA48/2 from π^0 decays. WASA-at-COSY has also published a bound based on π^0 decays, but it is weaker and applies only for $m_X > 20$ MeV [21]. Potentially more problematic is a bound from HADES, which searches for X bosons in π^0 , η , and Δ decays and excludes the dark photon parameter $\epsilon \gtrsim 3 \times 10^{-3}$, but this also applies only for $m_X > 20$ MeV [22]. Note also that $\pi^0 \rightarrow XX \rightarrow e^+e^-e^+e^-$ is not suppressed by the protophobic charge assignments, but it is suppressed by ϵ_n^4 and, for $|\epsilon_n| \sim 10^{-2}$, this is below current sensitivities. Similar considerations suppress X contributions to other decays, such as $\pi^+ \rightarrow \mu^+\nu_\mu e^+e^-$, to acceptable levels.

The hadronic charge can also be bounded by limits on Yukawa potentials from neutron-nucleus scattering. For a Yukawa potential $-g_n^2 A e^{-m_X r}/(4\pi r)$, n -Pb scattering requires $g_n^2/(4\pi) < 3.4 \times 10^{-11} (m_X/\text{MeV})^4$ [23]. The protophobic X boson induces a Yukawa potential $\epsilon_n^2 \alpha (A - Z) e^{-m_X r}/r$. Given $Z = 82$ and $A = 208$ for Pb, the bounds imply $|\epsilon_n| < 2.5 \times 10^{-2}$.

There are constraints from proton fixed target experiments. The ν -Cal I experiment at the U70 accelerator at IHEP provides a well-known dark photon constraint, but

its bounds are derived from X -bremsstrahlung from the initial p beam and π^0 decays to X bosons [24]. Both of these are suppressed in protophobic models. The CHARM experiment at CERN also bounds the parameter space through searches for $\eta, \eta' \rightarrow X\gamma$, followed by $X \rightarrow e^+e^-$ [25]. At the upper boundary of the region excluded by CHARM, the constraint is determined almost completely by the parameters that enter the X decay length, and so the dark photon bound on ε applies to ε_e and requires $|\varepsilon_e| > 2 \times 10^{-5}$. A similar, but weaker constraint can be derived from LSND data [26–28].

There are also bounds on the neutrino charge ε_ν . In the present case, where ε_e is non-zero, a recent study of $B-L$ gauge bosons [29] finds that these couplings are most stringently bounded by precision studies of $\bar{\nu} - e$ scattering from TEXONO for the m_X of interest here [30]. Reinterpreted for the present case, these studies require $|\varepsilon_\nu \varepsilon_e|^{1/2} \lesssim 7 \times 10^{-5}$. There are also bounds from coherent neutrino-nucleus scattering. Dark matter experiments with Xe target nuclei require a $B-L$ gauge boson to have coupling $g_{B-L} \lesssim 4 \times 10^{-5}$ [31]. Rescaling this to the current case, given $Z = 54$ and $A = 131$ for Xe, we find $|\varepsilon_\nu \varepsilon_n|^{1/2} < 2 \times 10^{-4}$.

To explain the ${}^8\text{Be}$ signal, ε_n must be significantly larger than ε_e . Nevertheless, the $\bar{\nu} - e$ scattering constraint provides a bound on ε_ν that is comparable to or stronger than the $\nu - N$ constraint throughout parameter space, and so we use the $\bar{\nu} - e$ constraint below. Note also that, given the range of acceptable ε_e , the bounds on ε_ν are more stringent than the bounds on ε_e , and so $B(X \rightarrow e^+e^-) \approx 100\%$, justifying our assumption above.

Although not our main concern, there are also bounds on second-generation couplings. For example, NA48/2 also derives bounds on $K^+ \rightarrow \pi^+ X$, followed by $X \rightarrow e^+e^-$ [10]. However, this branching ratio vanishes for massless X and is highly suppressed for low m_X . For $m_X = 17$ MeV, the bound on ε_n is not competitive with those discussed above [9, 11]. KLOE-2 also searches for $\phi \rightarrow \eta X$ followed by $X \rightarrow e^+e^-$ and excludes the dark photon parameter $\varepsilon \lesssim 7 \times 10^{-3}$ [32]. This is similar numerically to bounds discussed above, and the strange quark charge ε_s can be chosen to satisfy this constraint.

In summary, in the extreme protophobic case with $m_X \approx 17$ MeV, the charges are required to satisfy $|\varepsilon_n| < 2.5 \times 10^{-2}$ and $2 \times 10^{-4} < |\varepsilon_e| < 1.4 \times 10^{-3}$, and $|\varepsilon_\nu \varepsilon_e|^{1/2} \lesssim 7 \times 10^{-5}$. Combining these with Eqs. (5) and (7), we find that a protophobic gauge boson with first-generation charges

$$\begin{aligned} \varepsilon_u &= -\frac{1}{3}\varepsilon_n \approx \pm 3.7 \times 10^{-3} \\ \varepsilon_d &= \frac{2}{3}\varepsilon_n \approx \mp 7.4 \times 10^{-3} \\ 2 \times 10^{-4} &\lesssim |\varepsilon_e| \lesssim 1.4 \times 10^{-3} \\ |\varepsilon_\nu \varepsilon_e|^{1/2} &\lesssim 7 \times 10^{-5} \end{aligned} \quad (10)$$




FIG. 2. The ${}^8\text{Be}$ signal region, along with current constraints discussed in the text (gray) and projected sensitivities of future experiments in the (m_X, ε_e) plane. For the ${}^8\text{Be}$ signal, the other couplings are assumed to be in the ranges given in Eq. (10); for all other contours, the other couplings are those of a dark photon.

explains the ${}^8\text{Be}$ anomaly by ${}^8\text{Be}^* \rightarrow {}^8\text{Be} X$, followed by $X \rightarrow e^+e^-$, consistent with existing constraints. For $|\varepsilon_e|$ near the upper end of the allowed range in Eq. (10) and $|\varepsilon_\mu| \approx |\varepsilon_e|$, the X boson also solves the $(g-2)_\mu$ puzzle, reducing the current 3.6σ discrepancy to below 2σ [9].

Conclusions. We find evidence in the recent observation of a 6.8σ anomaly in the e^+e^- distribution of nuclear ${}^8\text{Be}$ decays for a new vector gauge boson. The new particle mediates a fifth force with a characteristic length scale of 12 fm. The requirements of the signal, along with the many constraints from other experiments that probe these low energy scales, constrain the mass and couplings of the boson to small ranges: its mass is $m_X \approx 17$ MeV, and it has milli-charged couplings to up and down quarks and electrons, but with relatively suppressed (and possibly vanishing) couplings to protons (and neutrinos) relative to neutrons. If its lepton couplings are approximately generation-independent, the 17 MeV vector boson may simultaneously explain the existing 3.6σ deviation from SM predictions in the anomalous magnetic moment of the muon. It is also interesting to note that couplings of this magnitude, albeit in an axial vector case, may resolve a 3.2σ excess in $\pi^0 \rightarrow e^+e^-$ decays [33, 34].

To confirm the ${}^8\text{Be}$ signal, the most direct approach would be to look for other nuclear states that decay to discrete gamma rays with energies above 17 MeV through M1 or E1 electromagnetic transitions. Unfortunately, the ${}^8\text{Be}$ system is quite special and, to our knowledge, the ${}^8\text{Be}^*$ and ${}^8\text{Be}^{*'}$ states yield the most energetic such gamma rays of all the nuclear states.

Nevertheless there are myriad opportunities to test and confirm this explanation, including re-analysis of old data sets, ongoing experiments, and many planned and future experiments, including DarkLight [35], HPS [36], LHCb [37], MESA [38], Mu3e [39], VEPP-3 [40], and possibly also SeaQuest [41] and SHiP [42]. The ^8Be signal region and expected sensitivities of these experiments are shown in Fig. 2. Further details about the existing constraints, prospects for the future, and UV completions of the model discussed here will be presented elsewhere [43].

Acknowledgments. We thank Attila J. Krasznahorkay and Alexandra Gade for helpful correspondence. The work of J.L.F., B.F., I.G., J.S., T.M.P.T., and P.T. is supported in part by NSF Grant No. PHY-1316792. The work of S.G. is supported in part by the DOE Office of Nuclear Physics under contract DE-FG02-96ER40989. J.L.F. is supported in part by a Guggenheim Foundation grant and in part by Simons Investigator Award #376204.

-
- [1] T. D. Lee and C.-N. Yang, “Conservation of Heavy Particles and Generalized Gauge Transformations,” *Phys. Rev.* **98** (1955) 1501.
- [2] A. Franklin, *The Rise and Fall of the Fifth Force: Discovery, Pursuit, and Justification in Modern Physics*. American Institute of Physics, New York, 1993.
- [3] A. Krasznahorkay *et al.*, “Observation of Anomalous Internal Pair Creation in Be8 : A Possible Indication of a Light, Neutral Boson,” *Phys. Rev. Lett.* **116** (2016) 042501, [arXiv:1504.01527 \[nucl-ex\]](#).
- [4] D. R. Tilley, J. H. Kelley, J. L. Godwin, D. J. Millener, J. E. Purcell, C. G. Sheu, and H. R. Weller, “Energy levels of light nuclei A=8,9,10,” *Nucl. Phys.* **A745** (2004) 155–362.
- [5] M. E. Rose, “Internal Pair Formation,” *Phys. Rev.* **76** (1949) 678–681. [Erratum: *Phys. Rev.* 78, 184 (1950)].
- [6] J. Gulys, T. J. Ketel, A. J. Krasznahorkay, M. Csatsl, L. Csige, Z. Gcsi, M. Hunyadi, A. Krasznahorkay, A. Vitz, and T. G. Tornyi, “A pair spectrometer for measuring multipolarities of energetic nuclear transitions,” *Nucl. Instrum. Meth.* **A808** (2016) 21–28, [arXiv:1504.00489 \[nucl-ex\]](#).
- [7] J. L. Hewett *et al.*, “Fundamental Physics at the Intensity Frontier,” [arXiv:1205.2671 \[hep-ex\]](#).
- [8] B. Döbrich, J. Jaekel, F. Kahlhoefer, A. Ringwald, and K. Schmidt-Hoberg, “ALPtraum: ALP production in proton beam dump experiments,” *JHEP* **02** (2016) 018, [arXiv:1512.03069 \[hep-ph\]](#).
- [9] M. Pospelov, “Secluded U(1) below the weak scale,” *Phys. Rev.* **D80** (2009) 095002, [arXiv:0811.1030 \[hep-ph\]](#).
- [10] **NA48/2** Collaboration, J. R. Batley *et al.*, “Search for the dark photon in π^0 decays,” *Phys. Lett.* **B746** (2015) 178–185, [arXiv:1504.00607 \[hep-ex\]](#).
- [11] H. Davoudiasl, H.-S. Lee, and W. J. Marciano, “Muon $g - 2$, rare kaon decays, and parity violation from dark bosons,” *Phys. Rev.* **D89** (2014) 095006, [arXiv:1402.3620 \[hep-ph\]](#).
- [12] **KLOE-2** Collaboration, A. Anastasi *et al.*, “Limit on the production of a low-mass vector boson in $e^+e^- \rightarrow U\gamma$, $U \rightarrow e^+e^-$ with the KLOE experiment,” *Phys. Lett.* **B750** (2015) 633–637, [arXiv:1509.00740 \[hep-ex\]](#).
- [13] **BaBar** Collaboration, J. P. Lees *et al.*, “Search for a Dark Photon in e^+e^- Collisions at BaBar,” *Phys. Rev. Lett.* **113** (2014) 201801, [arXiv:1406.2980 \[hep-ex\]](#).
- [14] R. Essig *et al.*, “Working Group Report: New Light Weakly Coupled Particles,” in *Community Summer Study 2013: Snowmass on the Mississippi (CSS2013) Minneapolis, MN, USA, July 29-August 6, 2013*. 2013. [arXiv:1311.0029 \[hep-ph\]](#).
- [15] E. M. Riordan *et al.*, “A Search for Short Lived Axions in an Electron Beam Dump Experiment,” *Phys. Rev. Lett.* **59** (1987) 755.
- [16] J. D. Bjorken, R. Essig, P. Schuster, and N. Toro, “New Fixed-Target Experiments to Search for Dark Gauge Forces,” *Phys. Rev.* **D80** (2009) 075018, [arXiv:0906.0580 \[hep-ph\]](#).
- [17] M. Davier and H. Nguyen Ngoc, “An Unambiguous Search for a Light Higgs Boson,” *Phys. Lett.* **B229** (1989) 150.
- [18] J. D. Bjorken, S. Ecklund, W. R. Nelson, A. Abashian, C. Church, B. Lu, L. W. Mo, T. A. Nunamaker, and P. Rassmann, “Search for Neutral Metastable Penetrating Particles Produced in the SLAC Beam Dump,” *Phys. Rev.* **D38** (1988) 3375.
- [19] M. D. Diamond and P. Schuster, “Searching for Light Dark Matter with the SLAC Millicharge Experiment,” *Phys. Rev. Lett.* **111** no. 22, (2013) 221803, [arXiv:1307.6861 \[hep-ph\]](#).
- [20] A. Bross, M. Crisler, S. H. Pordes, J. Volk, S. Errede, and J. Wrbanek, “A Search for Shortlived Particles Produced in an Electron Beam Dump,” *Phys. Rev. Lett.* **67** (1991) 2942–2945.
- [21] **WASA-at-COSY** Collaboration, P. Adlarson *et al.*, “Search for a dark photon in the $\pi^0 \rightarrow e^+e^-\gamma$ decay,” *Phys. Lett.* **B726** (2013) 187–193, [arXiv:1304.0671 \[hep-ex\]](#).
- [22] **HADES** Collaboration, G. Agakishiev *et al.*, “Searching a Dark Photon with HADES,” *Phys. Lett.* **B731** (2014) 265–271, [arXiv:1311.0216 \[hep-ex\]](#).
- [23] R. Barbieri and T. E. O. Ericson, “Evidence Against the Existence of a Low Mass Scalar Boson from Neutron-Nucleus Scattering,” *Phys. Lett.* **B57** (1975) 270–272.
- [24] J. Blümlein and J. Brunner, “New Exclusion Limits on Dark Gauge Forces from Proton Bremsstrahlung in Beam-Dump Data,” *Phys. Lett.* **B731** (2014) 320–326, [arXiv:1311.3870 \[hep-ph\]](#).
- [25] S. N. Gninenko, “Constraints on sub-GeV hidden sector gauge bosons from a search for heavy neutrino decays,” *Phys. Lett.* **B713** (2012) 244–248, [arXiv:1204.3583 \[hep-ph\]](#).
- [26] **LSND** Collaboration, C. Athanassopoulos *et al.*, “Evidence for muon-neutrino \rightarrow electron-neutrino oscillations from pion decay in flight neutrinos,” *Phys. Rev.* **C58** (1998) 2489–2511, [arXiv:nucl-ex/9706006 \[nucl-ex\]](#).
- [27] B. Batell, M. Pospelov, and A. Ritz, “Exploring Portals to a Hidden Sector Through Fixed Targets,” *Phys. Rev.* **D80** (2009) 095024, [arXiv:0906.5614 \[hep-ph\]](#).
- [28] R. Essig, R. Harnik, J. Kaplan, and N. Toro,

- “Discovering New Light States at Neutrino Experiments,” *Phys. Rev.* **D82** (2010) 113008, [arXiv:1008.0636 \[hep-ph\]](#).
- [29] S. Bilmis, I. Turan, T. M. Aliev, M. Deniz, L. Singh, and H. T. Wong, “Constraints on Dark Photon from Neutrino-Electron Scattering Experiments,” *Phys. Rev.* **D92** (2015) 033009, [arXiv:1502.07763 \[hep-ph\]](#).
- [30] **TEXONO** Collaboration, M. Deniz *et al.*, “Measurement of $\text{Nu}(e)\text{-bar}$ -Electron Scattering Cross-Section with a CsI(Tl) Scintillating Crystal Array at the Kuo-Sheng Nuclear Power Reactor,” *Phys. Rev.* **D81** (2010) 072001, [arXiv:0911.1597 \[hep-ex\]](#).
- [31] D. G. Cerdeo, M. Fairbairn, T. Jubbs, P. A. N. Machado, A. C. Vincent, and C. B. hm, “Physics from solar neutrinos in dark matter direct detection experiments,” [arXiv:1604.01025 \[hep-ph\]](#).
- [32] **KLOE-2** Collaboration, D. Babusci *et al.*, “Limit on the production of a light vector gauge boson in phi meson decays with the KLOE detector,” *Phys. Lett.* **B720** (2013) 111–115, [arXiv:1210.3927 \[hep-ex\]](#).
- [33] **KTeV** Collaboration, E. Abouzaid *et al.*, “Measurement of the rare decay $\pi^0 \rightarrow e^+e^-$,” *Phys. Rev.* **D75** (2007) 012004, [arXiv:hep-ex/0610072 \[hep-ex\]](#).
- [34] Y. Kahn, M. Schmitt, and T. M. P. Tait, “Enhanced rare pion decays from a model of MeV dark matter,” *Phys. Rev.* **D78** (2008) 115002, [arXiv:0712.0007 \[hep-ph\]](#).
- [35] J. Balewski *et al.*, “The DarkLight Experiment: A Precision Search for New Physics at Low Energies,” 2014. [arXiv:1412.4717 \[physics.ins-det\]](#).
- [36] O. Moreno, “The Heavy Photon Search Experiment at Jefferson Lab,” in *Meeting of the APS Division of Particles and Fields (DPF 2013) Santa Cruz, California, USA, August 13-17, 2013*. 2013. [arXiv:1310.2060 \[physics.ins-det\]](#).
- [37] P. Ilten, J. Thaler, M. Williams, and W. Xue, “Dark photons from charm mesons at LHCb,” *Phys. Rev.* **D92** (2015) 115017, [arXiv:1509.06765 \[hep-ph\]](#).
- [38] T. Beranek, H. Merkel, and M. Vanderhaeghen, “Theoretical framework to analyze searches for hidden light gauge bosons in electron scattering fixed target experiments,” *Phys. Rev.* **D88** (2013) 015032, [arXiv:1303.2540 \[hep-ph\]](#).
- [39] B. Echenard, R. Essig, and Y.-M. Zhong, “Projections for Dark Photon Searches at Mu3e,” *JHEP* **01** (2015) 113, [arXiv:1411.1770 \[hep-ph\]](#).
- [40] B. Wojtsekhowski, D. Nikolenko, and I. Rachek, “Searching for a new force at VEPP-3,” [arXiv:1207.5089 \[hep-ex\]](#).
- [41] S. Gardner, R. J. Holt, and A. S. Tadepalli, “New Prospects in Fixed Target Searches for Dark Forces with the SeaQuest Experiment at Fermilab,” [arXiv:1509.00050 \[hep-ph\]](#).
- [42] **SHiP** Collaboration, M. Anelli *et al.*, “A facility to Search for Hidden Particles (SHiP) at the CERN SPS,” [arXiv:1504.04956 \[physics.ins-det\]](#).
- [43] J. L. Feng, B. Fornal, I. Galon, S. Gardner, J. Smolinsky, T. M. P. Tait, and P. Tanedo. Work in progress.