

Squaring (1), averaging over spin, and performing the phase-space integration, we obtain for the total K -production cross section at threshold

$$\sigma_{p\Lambda K}(W) = \frac{1}{\sqrt{2}\pi} \left(\frac{G_{N\Lambda K}^2}{4\pi} \right) \frac{m_K^2}{M_\Lambda} \left(1 + \frac{M_\Lambda - M_N}{m_K} \right)^{-2} \times \left(\frac{1}{p^2 W} \right) (I_s + 2I_t), \quad (2a)$$

$$I_{s,t} = \int_0^\epsilon \eta \left(\frac{\epsilon - \eta}{m_K} \right)^{1/2} \sigma_{s,t}(\eta) d\eta, \quad (2b)$$

$$W = 2M + m_K + \epsilon = 2(p^2 + M_N^2)^{1/2} = 2M_N(1 + T_{\text{lab}}/2M_N)^{1/2}, \quad (2c)$$

where $\sigma_{s,t}(\eta)$ is the singlet (triplet) total Λp cross section at total c.m. kinetic energy η , and p is the c.m. momentum of the initial protons.⁶

⁶ Equation (2a) is formally different from Eq. (8a) of Ref. 1 because relativistic kinematics are used here; also, we include a factor, $4(\eta/m_K)^{1/2}$, erroneously omitted from Eq. (8b) of Ref. 1.

The inelasticity of the off-shell Λp scattering is accounted for in (2a) by a correction factor—viz., the ratio of the final to initial Λp c.m. momentum. This factor, which multiplies the elastic Λp cross section, approaches unity at the higher energies but is important near threshold.

The cross section is estimated using an effective-range approximation for $\sigma_{s,t}(\eta)$. Results, using effective-range parameters obtained from both hypernuclei data and Λp scattering data,⁷ are shown in Fig. 2. The lowest-energy K -production data known to us occurs at $P_{\text{lab}} = 2.807$ GeV/c,⁸ which is well outside the region of validity for this calculation. Further experimental work is needed in the region of $P_{\text{lab}} = 2.40$ GeV/c.

It is clear that the above analysis also applies to the process $p p \rightarrow p \Sigma K$; however, since the square of the $p \Sigma K$ coupling constant is an order of magnitude less than that for $p \Lambda K$,⁵ the predicted cross section near threshold is probably too small ($\sim 0.2 \mu\text{b}$) for experimental verification.

⁷ G. Alexander, O. Benary, U. Karshon, A. Shapira, G. Yekutieli, R. Engelmann, H. Filthuth, A. Fridman, and B. Schiby, Phys. Letters **19**, 715 (1966).

⁸ W. J. Fickinger, E. Pickup, D. K. Robinson, and E. O. Salant, Phys. Rev. **125**, 2082 (1962).

Asymptotic Sum Rules at Infinite Momentum*

J. D. BJORKEN

Stanford Linear Accelerator Center, Stanford University, Stanford, California

(Received 30 September 1968)

By combining the $q_0 \rightarrow i\infty$ method for asymptotic sum rules with the $P \rightarrow \infty$ method of Fubini and Furlan, we relate the structure functions W_2 and W_1 in inelastic lepton-nucleon scattering to matrix elements of commutators of currents at almost equal times at infinite momentum. We argue that the infinite-momentum limit for these commutators does not diverge, but may vanish. If the limit is nonvanishing, we predict $\nu W_2(\nu, q^2) \rightarrow f_2(\nu/q^2)$ and $W_1(\nu, q^2) \rightarrow f_1(\nu/q^2)$ as ν and q^2 tend to ∞ . From a similar analysis for neutrino processes, we conclude that at high energies the total neutrino-nucleon cross sections rise linearly with neutrino laboratory energy until nonlocality of the weak current-current coupling sets in. The sum of νp and $\bar{\nu} p$ cross sections is determined by the equal-time commutator of the Cabibbo current with its time derivative, taken between proton states at infinite momentum.

I. INTRODUCTION

INELASTIC lepton-nucleon scattering at high-momentum transfer is a very direct means of probing small-distance nucleon structure. Reflecting this fact is the profound state of theoretical ignorance on what, even qualitatively, can be expected in this process.¹ Some small inroads have been recently made using the

techniques of current algebra.² In particular, Cornwall and Norton³ have written down a large class of asymptotic sum rules, valid at large q^2 , for inelastic electron scattering. Of these, the sum rule of Callan and Gross⁴ relating an asymptotic integral over electron scattering cross sections to a piece of the commutator of electromagnetic current with its time derivative is of special interest. The purpose of this paper is to discuss such sum rules in a slightly different language—that of the infinite-

* Work supported by the U. S. Atomic Energy Commission.

¹ For a good example, see J. Bjorken, in *Proceedings of the International School of Physics "Enrico Fermi," Course 41*, edited by J. Steinberger (Academic Press Inc., New York, 1969); Stanford Linear Accelerator Center Report No. SLAC-PUB-338 (unpublished).

² J. D. Bjorken, Phys. Rev. **163**, 1767 (1967), and references quoted therein.

³ J. M. Cornwall and R. E. Norton, Phys. Rev. **177**, 2584 (1969).

⁴ C. G. Callan, Jr., and D. Gross, Phys. Rev. Letters **21**, 311 (1968).

momentum method. We show that the electron scattering data is related in a direct way to matrix elements of electromagnetic current commutators at infinite nucleon momentum.^{5,6} In particular, we find that the structure functions $W_2(q^2, \nu)$ and $W_1(q^2, \nu)$ describing inelastic scattering⁷ tend to simple limits for large q^2 :

$$\lim_{q^2 \rightarrow \infty, \nu/q^2 \text{ fixed}} \nu W_2(q^2, \nu) = M F_2(-q^2/M\nu), \quad (1.1)$$

$$\lim_{q^2 \rightarrow \infty, \nu/q^2 \text{ fixed}} M W_1(q^2, \nu) = F_1(-q^2/M\nu), \quad (1.2)$$

with

$$F_i(\omega) \equiv F_i(\omega) = \frac{-i}{\pi} \lim_{P_z \rightarrow \infty} \int_0^\infty d\tau \sin \omega \tau \int d^3x \times \langle P_z | [J_x(\mathbf{x}, \tau/P_0), J_x(0)] | P_z \rangle, \quad (1.3)$$

and

$$F_i(\omega) \equiv F_i(\omega) = \frac{F_2(\omega)}{\omega} - F_1(\omega) = \frac{i}{\pi} \lim_{P_z \rightarrow \infty} \int_0^\infty d\tau \sin \omega \tau \int d^3x \times \langle P_z | [J_z(\mathbf{x}, \tau/P_0), J_z(0)] | P_z \rangle, \quad (1.4)$$

where $\omega = -q^2/M\nu$. The existence of these limits (aside from the Brandt-Sucher disease⁸) is guaranteed by a finite value of the integral appearing in the Callan-Gross sum rule. Although the present data⁹ appear to indicate that F_2 is nonvanishing at $q^2 \sim 1-2 \text{ BeV}^2$, it is still possible that $F_2 \rightarrow 0$ and the infinite-momentum commutators in (1.3) and (1.4) vanish in the limit. In such a case the content of this paper is empty.

Sum rules such as Cornwall and Norton have written down⁸ may be obtained by taking the sine transform of (1.3) and (1.4) and expanding both sides in a power series in τ . For example, for $n=1, 3, 5, \dots$,

$$\begin{aligned} \int_0^\infty d\omega \omega^n F_i(\omega) &= \lim_{q^2 \rightarrow \infty} \left(\frac{|q^2|}{M} \right)^{n+1} \int_0^\infty \frac{d\nu}{\nu^{n+2}} W_1(q^2, \nu) \\ &= \lim_{P_z \rightarrow \infty} (-i)^{n/2} \int \frac{d^3x}{P_0^n} \\ &\quad \times \left\langle P_z \left| \left[\frac{\partial^n J_x(\mathbf{x}, t)}{\partial t^n}, J_x(0) \right] \right| P_z \right\rangle_{t=0} \\ &\quad n=1, 3, 5, \dots \end{aligned} \quad (1.5)$$

⁵ S. Fubini and G. Furlan, *Physics* **1**, 229 (1965).

⁶ R. Dashen and M. Gell-Mann, in *Proceedings of the Third Coral Gables Conference on Symmetry Principles at High Energy*, edited by B. Kursunoglu, A. Perlmutter, and I. Sakmar (W. H. Freeman and Co., San Francisco, 1966).

⁷ See, e.g., T. de Forest and J. Walecka, *Advan. Phys.* **15**, 1 (1966); and Eq. (2.2).

⁸ R. Brandt and J. Sucher, *Phys. Rev. Letters* **20**, 1131 (1968); and (to be published).

⁹ E. Bloom, D. Coward, H. DeStaebl, J. Drees, J. Litt, G. Miller, L. Mo, R. Taylor, M. Breidenbach, J. Friedman, G. Hartman, H. Kendall, and S. Loken, report to Fourteenth International Conference on High Energy Physics Vienna, 1968 (unpublished) and numerous private communications which we gratefully acknowledge.

with a similar expression for F_i or W_2 . However, the content of these results is more succinctly discussed in terms of (1.1)–(1.4).

Although a straightforward generalization of these relations to different currents and momentum states is not difficult, what is not straightforward is the interpretation of the almost equal-time commutators appearing in (1.3) and (1.4). In particular, the spectrum of intermediate “frequencies” $\omega = -q^2/M\nu$ is bounded above, corresponding to at most the intermediate energy appropriate to the single-nucleon Z diagram (see Fig. 1). Assuming that the limit (1.1) and (1.2) is nontrivial (nonvanishing), it will be most interesting to construct models with the kind of asymptotic behavior expressed in (1.1)–(1.4). This, however, is beyond the scope of this paper.

In Sec. II, a simple derivation of the result is given. Section III remedies the swindle perpetrated on the reader in Sec. II, by providing a more honest derivation. In Sec. IV, we attempt a generalization to an arbitrary kinematical situation. In Sec. V, we apply the same method to neutrino reactions and find that νp and $\bar{\nu} p$ total cross sections should rise linearly with energy. The sum of νp and $\bar{\nu} p$ cross sections is determined by the equal-time commutator of the Cabibbo current with its first time derivative. Section VI summarizes our conclusions.

II. SIMPLE DERIVATION OF THE ASYMPTOTIC LIMIT


The inelastic scattering cross section from an unpolarized nucleon may be written⁷ as

$$\begin{aligned} \frac{d\sigma}{d\Omega dE'} &= \frac{\alpha^2}{4E^2 \sin^4(\frac{1}{2}\theta)} \\ &\quad \times [W_2(q^2, \nu) \cos^2(\frac{1}{2}\theta) + 2W_1(q^2, \nu) \sin^2(\frac{1}{2}\theta)], \end{aligned} \quad (2.1)$$

where E, E' is the energy of incident and scattered electron, θ is the scattering angle of electron, q^2 equals $-4EE' \sin^2(\frac{1}{2}\theta)$, ν equals $q \cdot P/M = (E - E')$, P is the momentum of target nucleon, q is the momentum of virtual photon, and

$$\begin{aligned} &\frac{1}{M^2} \left(P_\mu - \frac{P \cdot q q_\mu}{q^2} \right) \left(P_\nu - \frac{P \cdot q q_\nu}{q^2} \right) W_2(q^2, \nu) \\ &\quad - \left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) W_1(q^2, \nu) \\ &= \frac{P_0}{M^n} \sum \langle P | J_\mu(0) | n \rangle \langle n | J_\nu(0) | P \rangle (2\pi)^3 \delta^4(P_n - P - q) \\ &= \frac{P_0}{M} \int \frac{d^4x}{2\pi} e^{iq \cdot x} \langle P | [J_\mu(x), J_\nu(0)] | P \rangle. \end{aligned} \quad (2.2)$$

Fig. 1. Single-nucleon
Z diagram.



In all matrix elements $\langle P | \dots | P \rangle$, an average over nucleon spin is implied. Now consider the limit of (2.2) as $P_0 \rightarrow \infty$, $q_0 \rightarrow \infty$, $q_0/P_0 \rightarrow -\omega$ fixed, \mathbf{q} fixed ($\omega = -q^2/M\nu$). Notice that $q^2 \rightarrow +\infty$ (timelike) in this limit.

Choosing $\mu \neq 0$, $\nu \neq 0$, we find

$$\frac{P_i P_j}{M^2} W_2 + \delta_{ij} W_1 \xrightarrow{P_z \rightarrow \infty} \frac{P_0}{M} \int d^3x \int_{-\infty}^{\infty} \frac{dt}{2\pi} \times e^{-i\omega P_0 t - i\mathbf{q} \cdot \mathbf{x}} \langle P | [J_i(\mathbf{x}, P_0 t/P_0), J_j(0)] | P \rangle, \quad (2.3)$$

or, using (1.1) and (1.2) and the assumption that the commutator vanishes outside the light cone,

$$\begin{aligned} & -\delta_{i3}\delta_{j3} \frac{F_2(\omega)}{\omega} + \delta_{ij} F_1(\omega) \\ &= \lim_{P_z \rightarrow \infty} \int d^3x \int_{-\infty}^{\infty} \frac{d\tau}{2\pi} e^{-i\omega\tau} \langle P | [J_i(\mathbf{x}, \tau/P_0), J_j(0)] | P \rangle \\ &= -\frac{i}{\pi} \lim_{P_z \rightarrow \infty} \int d^3x \int_0^{\infty} d\tau \sin \omega\tau \\ & \quad \times \langle P | [J_i(\mathbf{x}, \tau/P_0), J_j(0)] | P \rangle \quad (\omega < 0). \quad (2.4) \end{aligned}$$

Now (2.4) defines F_1 and F_2 for positive ω as well as negative; therefore we let $\omega = +|q^2|/M\nu > 0$, as appropriate for inelastic scattering. Then we get

$$\begin{aligned} F_i(\omega) &= F_1(\omega) = -\frac{i}{\pi} \lim_{P_z \rightarrow \infty} \int d^3x \int_0^{\infty} d\tau \sin |\omega| \tau \\ & \quad \times \langle P | [J_i(\mathbf{x}, \tau/P_0), J_x(0)] | P \rangle \quad (\omega > 0) \quad (2.5) \end{aligned}$$

and

$$\begin{aligned} F_i(\omega) &\equiv \frac{F_2(\omega)}{|\omega|} - F_1(\omega) = +\frac{i}{\pi} \lim_{P_z \rightarrow \infty} \int d^3x \int_0^{\infty} d\tau \sin |\omega| \tau \\ & \quad \times \langle P | [J_z(\mathbf{x}, \tau/P_0), J_z(0)] | P \rangle \quad (\omega > 0). \quad (2.6) \end{aligned}$$

This is the desired result given in (1.3) and (1.4). Both F_1 and F_2 are positive; notice the curious sign change between the transverse and longitudinal commutators.

The reader should have noticed the swindle that has been perpetrated in letting $\omega \rightarrow -\omega$. There has been no justification that $\omega < 0$ can be extrapolated from $\omega > 0$. Section III is devoted to providing such a justification.

III. JUSTIFICATION OF THE RESULTS

In order to provide a better derivation of the preceding results, we consider the covariant current correlation

function

$$\begin{aligned} T_{\mu\nu}^* &= \frac{1}{M^2} \left(P_\mu - \frac{P \cdot q q_\mu}{q^2} \right) \left(P_\nu - \frac{P \cdot q q_\nu}{q^2} \right) T_2(q^2, \nu) \\ & \quad - \left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) T_1(q^2, \nu) \\ &= +\frac{iP_0}{M} \int d^4x e^{iq \cdot x} \theta(x_0) \langle P | [J_\mu(x), J_\nu(0)] | P \rangle \\ & \quad + \text{polynomials in } q, \quad (3.1) \end{aligned}$$

where, as always in this paper, an average over nucleon spin is implied. According to the dictum of Harari,¹⁰ T_2 satisfies an unsubtracted dispersion relation in ν , while T_1 requires one subtraction, provided $q^2 < 0$, i.e., spacelike.

$$T_2 = \frac{1}{\pi} \int_0^\infty \frac{d\nu'^2 \text{Im} T_2(\nu', q^2)}{\nu'^2 - \nu^2 - i\epsilon}, \quad (3.2)$$

$$\begin{aligned} T_1(\nu, q^2) &= T_1(0, q^2) \\ & \quad + \frac{\nu^2}{\pi} \int_0^\infty \frac{d\nu'^2 \text{Im} T_1(\nu', q^2)}{\nu'^2(\nu'^2 - \nu^2 - i\epsilon)} \quad (q^2 < 0). \quad (3.3) \end{aligned}$$

We choose to express the T_i in terms of the variables q^2 and $\omega = (-q^2/M\nu)$. Using the fact that

$$\frac{1}{\pi} \text{Im} T_i(\nu, q^2) = W_i(\nu, q^2), \quad (3.4)$$

we obtain

$$T_1(\omega, q^2) = T_1(\infty, q^2) - \int_0^4 \frac{d\omega'^2 W_1(\omega', q^2)}{(\omega'^2 - \omega^2 + i\epsilon)}, \quad (3.5)$$

$$T_2(\omega, q^2) = -\omega^2 \int_0^4 \frac{d\omega'^2 W_2(\omega', q^2)}{\omega'^2(\omega'^2 - \omega^2 + i\epsilon)}. \quad (3.6)$$

We now take the limit $q_0 \rightarrow i\infty$, \mathbf{q} fixed and P_z temporarily fixed. In this limit

$$\omega = \frac{-q^2}{q_0 P_0} \rightarrow -\frac{i|q_0|}{P_0} \rightarrow -i\infty,$$

$$T_1 \rightarrow T_1(\infty, q^2) - \frac{P_0^2}{|q_0|^2} \int_0^4 d\omega'^2 W_1(\omega', q^2),$$

$$T_2 \rightarrow + \int_0^4 \frac{d\omega'^2}{\omega'^2} W_2(\omega', q^2).$$

¹⁰ H. Harari, Phys. Rev. Letters **17**, 1303 (1963).

On the other hand, from (3.1),

$$\begin{aligned} T_{\mu\nu} &\rightarrow \frac{iP_0}{M} \int d^3x \int_0^\infty dt e^{-i q_0 t} \langle P | [J_\mu(x), J_\nu(0)] | P \rangle \\ &\quad + \text{polynomial} \\ &\rightarrow \frac{iP_0}{M |q_0|} \int d^3x \langle P | [J_\mu(\mathbf{x}, 0), J_\nu(0)] | P \rangle + \frac{iP_0}{M |q_0|^2} \\ &\quad \times \int d^3x \langle P | [\partial J_\mu(\mathbf{x}, t) / \partial t, J_\nu(0)] | P \rangle_{t=0} + \dots \quad (3.7) \\ &\quad + \text{polynomial} \end{aligned}$$

Specializing to μ and $\nu \neq 0$, and \mathbf{P} in the z direction, we find

$$\begin{aligned} \frac{P_0}{M} \int d^3x \langle P | [\partial J_z(\mathbf{x}, t) / \partial t, J_x(0)] | P \rangle_{t=0} + \text{polynomial} \\ = \lim_{q^2 \rightarrow -\infty} i q^2 T_1(\infty, q^2) + i P_0^2 \int_0^4 d\omega'^2 W_1(\omega', q^2) \quad (3.8) \\ \frac{P_0}{M} \int d^3x \langle P | [\partial J_z(\mathbf{x}, t) / \partial t, J_z(0)] | P \rangle_{t=0} \\ - \frac{P_0}{M} \int d^3x \left\langle P \left| \left[\frac{\partial J_z(\mathbf{x}, t)}{\partial t}, J_x(0) \right] \right| P \right\rangle_{t=0} \\ = \lim_{q^2 \rightarrow -\infty} i q^2 \frac{P_z^2}{M^2} \int_0^4 \frac{d\omega'^2}{\omega'^2} W_2(\omega', q^2). \quad (3.9) \end{aligned}$$

Hereafter, we shall assume that

$$\begin{aligned} \lim_{q^2 \rightarrow -\infty} |q^2| \int_0^4 \frac{d\omega'^2}{\omega'^2} W_2(\omega', q^2) \\ = \lim_{q^2 \rightarrow -\infty} 2 |q^2| \int_0^\infty \frac{d\nu'}{\nu'} W_2(\nu', q^2) < \infty. \quad (3.10) \end{aligned}$$

For fixed q^2 , the integration in ν converges, if the Harari dictum¹⁰ is correct. The $|q^2| \rightarrow \infty$ limit is that taken by Callan and Gross²; in particular, (3.10) is an integral involved in their sum rules. It is unlikely the inelastic scattering is so large that this Callan-Gross integral does not exist; such a circumstance would oversaturate the sum rules for $\int d\nu W_2$ from current algebra.¹ More likely is the vanishing of (3.10). If the Callan-Gross integral exists, i.e., (3.10) holds, we can show that almost-equal-time commutators (1.3) and (1.4) necessarily exist as well. To show this, we observe from their kinematical definitions in terms of transverse and longitudinal cross sections¹¹ that in our limit

$$\frac{W_1}{W_2} = \left(1 + \frac{\nu^2}{|q^2|}\right) \left(\frac{\sigma_t}{\sigma_t + \sigma_l}\right) \leq \frac{\nu^2}{|q|^2}, \quad (3.11)$$

¹¹ For a definition of σ_t and σ_l , see L. Hand, in *Proceedings of the 1967 International Symposium on Electron and Photon Interactions at High Energies* (Stanford Linear Accelerator Center, Stanford,

which implies

$$q^4 \int \frac{d\nu'}{\nu'^3} W_1 \leq |q^2| \int \frac{d\nu'}{\nu'} W_2 < \infty. \quad (3.12)$$

In other words,

$$\lim_{q^2 \rightarrow -\infty} \int_0^4 d\omega'^2 W_1(\omega', q^2) < \infty, \quad (3.13)$$

$$\lim_{q^2 \rightarrow -\infty} |q^2| \int \frac{d\omega'^2}{\omega'^2} W_2(\omega', q^2) < \infty, \quad (3.14)$$

and because W_1 and W_2 are positive semidefinite, it follows that for $|\omega| > 2$,

$$\phi_1(\omega) = \lim_{q^2 \rightarrow -\infty} \int_0^4 \frac{d\omega'^2 W_1(\omega', q^2)}{\omega^2 - \omega'^2} < \infty, \quad (3.15)$$

and ϕ therefore by analytic continuation exists throughout the cut ω plane, barring extreme pathology in the behavior of W_1 in the limit. Similarly, we have

$$\phi_2(\omega) = \lim_{q^2 \rightarrow -\infty} |q^2| \int_0^4 \frac{d\omega'^2 W_2(\omega', q^2)}{\omega'^2(\omega^2 - \omega'^2)} < \infty \quad (3.16)$$

throughout the cut ω plane. These results are then sufficient to guarantee the existence of F_1 , F_2 , and the almost-equal-time commutators (1.3) and (1.4). We go back to $T_{\mu\nu}^*$ as defined in (3.1), (3.5), and (3.6) and let $P_z \rightarrow \infty$, $q_0 \rightarrow i\infty$, \mathbf{q} fixed, $\omega = -q_0/P_0 = -i|q_0/P_0|$ fixed.

$$\begin{aligned} T_{ij}^* &\rightarrow + \frac{P_i P_j}{M^2} |\omega|^2 \int_0^4 \frac{d\omega'^2 W_2(\omega', q^2)}{\omega'^2(\omega'^2 + |\omega|^2)} \\ &\quad + \delta_{ij} \left[T_1(\infty, q^2) - \int_0^4 \frac{d\omega'^2 W_1(\omega', q^2)}{(\omega'^2 + |\omega|^2)} \right]. \quad (3.17) \end{aligned}$$

In terms of F_1 and F_2 , we find, from (1.1), (1.2), and (3.1),

$$\begin{aligned} T_{ij}^* &\rightarrow \frac{P_i P_j}{|q^2|} \frac{|\omega|^2}{M} \int_0^4 \frac{d\omega'^2 F_2(\omega')}{|\omega'|(\omega'^2 + |\omega|^2)} \\ &\quad + \delta_{ij} \left[T_1(\infty, q^2) - \frac{1}{M} \int_0^4 \frac{d\omega'^2 F_1(\omega')}{\omega'^2 + |\omega|^2} \right] \\ &= \lim_{P_z \rightarrow \infty} \frac{i}{M} \int d^3x \int_0^\infty d\tau e^{-i|\omega|\tau} \langle P | [J_i(\mathbf{x}, \tau/E), J_j(0)] | P \rangle \\ &\quad + \text{polynomial}. \quad (3.18) \end{aligned}$$

Calif., 1967). The connection between W_1 , W_2 and σ_t , σ_l is

$$W_1 = \frac{\sigma_t}{4\pi^2\alpha} \left(\nu - \frac{|q^2|}{2M} \right), \quad W_2 = \frac{(\sigma_t + \sigma_l)(\nu - q^2/2M)}{4\pi^2\alpha(1 + \nu^2/|q^2|)}.$$

Notice that F_t and F_l , Eqs. (1.3) and (1.4), are proportional to σ_t and σ_l in the limit: $4\pi^2\alpha F_t/\nu \rightarrow (1 - \frac{1}{2}\omega)\sigma_t$, $4\pi^2\alpha F_l/\nu \rightarrow (1 - \frac{1}{2}\omega)\sigma_l$.

In the limit,

$$\frac{P_i P_j}{|q^2|} \rightarrow \delta_{i3} \delta_{j3} \left| \frac{P_0^2}{q_0^2} \right| \rightarrow \frac{\delta_{i3} \delta_{j3}}{|\omega^2|}$$

and

$$T_1(\infty, q^2) \rightarrow \text{polynomial.} \quad (3.19)$$

The existence of the commutator in (3.18) is guaranteed by the existence of an inverse Laplace transform of (3.18). Having taken the limit $|q^2| \rightarrow \infty$, etc., we may continue (3.18) into the cut ω plane, and obtain

$$\begin{aligned} & \delta_{i3} \delta_{j3} \int_0^4 \frac{d\omega'^2 F_2(\omega')}{\omega'(\omega'^2 - \omega^2 - i\epsilon)} - \delta_{ij} \int_0^4 \frac{d\omega'^2 F_1(\omega')}{\omega'^2 - \omega^2 - i\epsilon} \\ &= \lim_{P_z \rightarrow \infty} i \int d^3x \int_0^\infty d\tau \\ & \quad \times e^{i\omega\tau} \langle P | [J_i(x, \tau/P_0), J_j(0)] | P \rangle. \end{aligned} \quad (3.20)$$

Upon taking the imaginary part of this relation, we reproduce (2.5) and (2.6). This justifies the short derivation given in Sec. II.

IV. GENERALIZATION

The preceding analysis can be generalized to arbitrary currents and momenta of the states. As an example, we

consider the case of two different $SU(3) \times SU(3)$ currents (q_1, q_2) sandwiched between spin-zero hadron states (p_1, p_2) of the same parity. We use the notation of Bander and Bjorken¹²:

$$\begin{aligned} p_1 + q_1 &\rightarrow p_2 + q_2, \\ P &= p_1 + p_2, \\ \Delta &= p_2 - p_1 = q_1 - q_2, \\ Q &= q_1 + q_2, \end{aligned} \quad (4.1)$$

$$\nu = P \cdot Q, \quad t = \Delta^2, \quad \delta = \Delta \cdot Q = q_1^2 - q_2^2,$$

and take the limit

$$E \simeq p_{1z} = p_{2z} \rightarrow \infty, \quad Q_0 \rightarrow i\infty, \quad \mathbf{Q} = 0 \quad (4.2)$$

such that

$$\omega = -Q_0/P_0 \simeq -Q^2/\nu \quad (4.3)$$

remains finite. Encouraged by the reasonableness of this limit in the special case of Secs. I–III, we *assume* that it exists in this case as well.

Under these circumstances, the new general invariants δ and t tend to a finite limit:

$$\begin{aligned} \delta &= Q \cdot \Delta \rightarrow -\omega[(p_{21}^2 + m_2^2) - (p_{11}^2 + m_1^2)], \\ t &= \Delta^2 \rightarrow -(\mathbf{p}_{21} - \mathbf{p}_{11})^2. \end{aligned} \quad (4.4)$$

The covariant amplitude $M_{\mu\nu}^{\alpha\beta*}$ tends, in the limit, to

$$\begin{aligned} M_{\mu\nu}^{\alpha\beta*} &= -(2\pi)^3 i (4\omega_1 \omega_2)^{1/2} \int d^4x e^{iq_1 \cdot x} \theta(x_0) \langle P_2 | [j_\mu^\alpha(x), j_\nu^\beta(0)] | P_1 \rangle \\ &\rightarrow -(2\pi)^3 i (2E) \int d^3x \int_0^\infty dt e^{-(|Q_0|t/2)} \lim_{P_z \rightarrow \infty} \langle P_z, \mathbf{P}_{21} | [j_\mu^\alpha(x, t), j_\nu^\beta(0)] | P_z, \mathbf{P}_{11} \rangle \\ &\rightarrow -2i(2\pi)^3 \int d^3x \int_0^\infty d\tau e^{-|\omega|\tau} \lim_{P_z \rightarrow \infty} \langle P_z, \mathbf{P}_{21} | [j_\mu^\alpha(x, \tau/E), j_\nu^\beta(0)] | P_z, \mathbf{P}_{11} \rangle (+\text{polynomials}). \end{aligned} \quad (4.5)$$

This last expression is a function of \mathbf{p}_{21} , \mathbf{p}_{11} , and ω alone. Upon writing out $M_{\mu\nu}^{\alpha\beta}$ in invariants (suppressing indices $\alpha\beta$)

$$\begin{aligned} M_{\mu\nu} &= P_\mu P_\nu A_1 + (P_\mu Q_\nu + P_\nu Q_\mu) A_2 + (P_\mu Q_\nu - P_\nu Q_\mu) A_3 \\ &+ (P_\mu \Delta_\nu + P_\nu \Delta_\mu) A_4 + (P_\mu \Delta_\nu - P_\nu \Delta_\mu) A_5 \\ &+ (Q_\mu \Delta_\nu + Q_\nu \Delta_\mu) A_6 + (Q_\mu \Delta_\nu - Q_\nu \Delta_\mu) A_7 \\ &+ Q_\mu Q_\nu A_8 + \Delta_\mu \Delta_\nu A_9 + g_{\mu\nu} A_{10}, \end{aligned} \quad (4.6)$$

we see that A_4 , A_5 , A_6 , and A_7 would have to tend to $(Q^2)^{-1/2}$ in order that the limit be nonvanishing and finite. We consider this unlikely, but cannot exclude it. Here we *assume* that in the limit these A_i do not contribute.

We write

$$\begin{aligned} A_i^{\alpha\beta} &\rightarrow \frac{1}{Q^2} F_i^{\alpha\beta}(\omega, t, \epsilon), \quad i=1, 2, 3, 8 \\ Q_0 A_i^{\alpha\beta} &\rightarrow 0, \quad i=4, 5, 6, 7 \\ A_i^{\alpha\beta} &\rightarrow F_i^{\alpha\beta}(\omega, t, \epsilon), \quad i=9, 10 \end{aligned} \quad (4.7)$$

where we introduce the variable

$$\epsilon = -\delta/\omega = (p_{21}^2 + m_2^2) - (p_{11}^2 + m_1^2). \quad (4.8)$$

When these limits (4.7) are inserted into (4.6), we find in the asymptotic infinite-momentum limit

$$\begin{aligned} M_{\mu\nu}^{\alpha\beta} &\rightarrow \frac{\theta_\mu \theta_\nu}{\omega^2} F_1 - \frac{(\theta_\mu \eta_\nu + \theta_\nu \eta_\mu)}{\omega} F_2 - \frac{(\theta_\mu \eta_\nu - \theta_\nu \eta_\mu)}{\omega} F_3 \\ &+ \eta_\mu \eta_\nu F_8 + \Delta_\mu \Delta_\nu F_9 + g_{\mu\nu} F_{10}, \end{aligned} \quad (4.9)$$

where $\theta_\mu = (1, 1, 0, 0)$, $\eta_\mu = (1, 0, 0, 0)$.

We are now free to identify various combinations of these form factors in terms of the almost-equal-time current commutators at infinite momentum. Using i or

¹² M. Bander and J. Bjorken, Phys. Rev. **174**, 1704 (1968).

j to indicate transverse components,

$$\begin{aligned}
 M_{00}^{\alpha\beta} &\rightarrow \frac{F_1^{\alpha\beta}}{\omega^2} - \frac{2F_2^{\alpha\beta}}{\omega} + F_8^{\alpha\beta} + F_{10}^{\alpha\beta}, & M_{zz}^{\alpha\beta} &\rightarrow \frac{F_1^{\alpha\beta}}{\omega^2} - F_{10}^{\alpha\beta}, \\
 M_{0z}^{\alpha\beta} &\rightarrow \frac{-F_2^{\alpha\beta} + F_3^{\alpha\beta}}{\omega}, & M_{iz}^{\alpha\beta} &\rightarrow 0, \\
 M_{z0}^{\alpha\beta} &\rightarrow \frac{-F_2^{\alpha\beta} - F_3^{\alpha\beta}}{\omega}, & M_{ij}^{\alpha\beta} &\rightarrow (\Delta_i \Delta_j F_9^{\alpha\beta} - \delta_{ij} F_{10}^{\alpha\beta}), \\
 M_{0i}^{\alpha\beta} &\rightarrow 0, & M_{i0}^{\alpha\beta} &\rightarrow 0.
 \end{aligned} \tag{4.10}$$

Recall

$$M_{\mu\nu}^{\alpha\beta} \rightarrow -2i(2\pi)^3 \int d^3x \int_0^\infty d\tau e^{-i\omega|\tau|} \langle \mathbf{P}_{21} | [j_\mu^\alpha(\mathbf{x}, \tau/E), j_\nu^\beta(0)] | \mathbf{P}_{11} \rangle_{P_z \rightarrow \infty} \tag{4.11}$$

(+polynomials)

Thus all invariant functions F_i can be determined in terms of the various current correlation functions, which then play the central role. Similarly, an infinite set of convergent sum rules, whose right-hand side involves commutators of J_ν with $\partial_0^n J_\mu$, can be obtained by expanding (4.10) and (4.11) in inverse powers of ω , and comparing coefficients as $\omega \rightarrow \infty$. These are independent of the asymptotic sum rules of Bander and Bjorken,¹² because in this case δ does not tend to zero, but rather to ∞ . We do not know what to do with these results, and shall not pursue them further here.

V. NEUTRINO PROCESSES

If we write the analog of (2.1) for the process $\bar{\nu}_\mu + P \rightarrow \mu^+ + \text{hadrons}$ as^{1,13,14}

$$\begin{aligned}
 \frac{\pi}{EE'} \frac{d\sigma}{d\Omega dE'} &= \frac{M d\sigma}{d|q^2| d\nu} \\
 &= \frac{E'}{E} \frac{G^2}{2\pi} \left[\mathbf{W}_2 \cos^2(\tfrac{1}{2}\theta) + 2\mathbf{W}_1 \sin^2(\tfrac{1}{2}\theta) \right. \\
 &\quad \left. + \frac{(E+E')}{M} \mathbf{W}_3 \sin^2(\tfrac{1}{2}\theta) \right] \tag{5.1}
 \end{aligned}$$

with kinematics as in Sec. II, and

$$\begin{aligned}
 \frac{P_0}{M} \int \frac{d^4x}{2\pi} e^{iq \cdot x} \langle P | [j_\mu(x), j_\nu^\dagger(0)] | P \rangle \\
 = \frac{P_\mu P_\nu}{M^2} \mathbf{W}_2 - g_{\mu\nu} \mathbf{W}_1 - i \frac{\epsilon_{\mu\nu\alpha\beta} P^\alpha q^\beta}{2M^2} \mathbf{W}_3 + \dots, \tag{5.2}
 \end{aligned}$$

where j_μ is the Cabibbo current,¹⁵ it follows from the

arguments in Sec. IV that under our assumptions

$$\begin{aligned}
 \mathbf{W}_2 &\rightarrow \frac{1}{\nu} \mathbf{F}_2 \left(\frac{-q^2}{M\nu} \right), & M\mathbf{W}_1 &\rightarrow \mathbf{F}_1 \left(\frac{-q^2}{M\nu} \right), \\
 \mathbf{W}_3 &\rightarrow \frac{1}{\nu} \mathbf{F}_3 \left(\frac{-q^2}{M\nu} \right),
 \end{aligned} \tag{5.3}$$

as $q^2 \rightarrow -\infty$, $\nu \rightarrow \infty$, $q^2/\nu \rightarrow \text{constant}$.

Introducing the variables $\omega = -q^2/M\nu$ and $x = \nu/E$, the total cross section coming from (5.1) becomes (for $E \gg M$)

$$\begin{aligned}
 \sigma_{\text{tot}} &\rightarrow \int_0^2 d\omega \int_0^1 dx \left(\frac{G^2 M E}{2\pi} \right) \\
 &\times [(1-x)\mathbf{F}_2(\omega) + \tfrac{1}{2}x^2\omega\mathbf{F}_1(\omega) + \tfrac{1}{2}x(1-\tfrac{1}{2}x)\omega\mathbf{F}_3(\omega)] \\
 &= \frac{G^2 M E}{4\pi} \int_0^2 d\omega [\mathbf{F}_2(\omega) + \tfrac{1}{3}\omega\mathbf{F}_1(\omega) + \tfrac{1}{3}\omega\mathbf{F}_3(\omega)]. \tag{5.4}
 \end{aligned}$$

Therefore, the cross section is predicted to rise linearly with laboratory neutrino energy. The coefficient is controlled again by the behavior of the current commutators at almost equal time and at infinite momentum. To determine this, we take various components of (5.2) in the $q_0, P_z \rightarrow \infty$ limit, in parallel with the discussion leading to (2.5):

$$\begin{aligned}
 \lim_{P_z \rightarrow \infty} \int d^3x \int_{-\infty}^\infty \frac{d\tau}{2\pi} \\
 \times e^{-i\omega\tau} \langle P | [j_x(\mathbf{x}, \tau/E), j_x^\dagger(0)] | P \rangle &= \mathbf{F}_1(\omega), \\
 \lim_{P_z \rightarrow \infty} \int d^3x \int_{-\infty}^\infty \frac{d\tau}{2\pi} \\
 \times e^{-i\omega\tau} \langle P | [j_z(\mathbf{x}, \tau/E), j_z^\dagger(0)] | P \rangle &= \left[\mathbf{F}_1(\omega) - \frac{\mathbf{F}_2(\omega)}{\omega} \right], \\
 \lim_{P_z \rightarrow \infty} \int d^3x \int_{-\infty}^\infty \frac{d\tau}{2\pi} \\
 \times e^{-i\omega\tau} \langle P | [j_x(\mathbf{x}, \tau/E), j_y^\dagger(0)] | P \rangle &= -\tfrac{1}{2}i\mathbf{F}_3(\omega). \tag{5.5}
 \end{aligned}$$

¹³ T. D. Lee and C. N. Yang, Phys. Rev. **126**, 2239 (1962).

¹⁴ S. Adler, Phys. Rev. **143**, 1144 (1966).

¹⁵ Our normalization is $j_\mu = \bar{\psi}' \gamma_\mu (1 - \gamma_5) (\psi' \cos\theta + \lambda' \sin\theta)$.

Substituting (5.5) into (5.4) we find, upon extending the ω integration to ∞ ,

$$\sigma_{\text{tot}} \rightarrow \frac{G^2 M E}{4\pi} \int_0^\infty d\omega \omega \int_{-\infty}^\infty \frac{d\tau}{2\pi} e^{-i\omega\tau} C(\tau), \quad (5.6)$$

where

$$C(\tau) = \lim_{P_z \rightarrow \infty} \int d^3x \langle P_z | \frac{4}{3} [j_x(\mathbf{x}, \tau/E), j_x^\dagger(0)] - [j_z(\mathbf{x}, \tau/E), j_z^\dagger(0)] + \frac{2}{3} i [j_x(\mathbf{x}, \tau/E), j_y^\dagger(0)] | P_z \rangle. \quad (5.7)$$

An interesting result is obtained upon taking the sum of antineutrino and neutrino cross sections. By crossing symmetry,

$$\begin{aligned} \sigma_{\text{tot}}^{\bar{\nu}p} + \sigma_{\text{tot}}^{\nu p} &= \frac{G^2 M E}{4\pi} \int_{-\infty}^\infty \omega d\omega \int_{-\infty}^\infty \frac{d\tau}{2\pi} e^{-i\omega\tau} C(\tau) = \frac{G^2 M E}{4\pi} (-i) \frac{\partial C(\tau)}{\partial \tau} \Big|_{\tau=0} \\ &= \frac{G^2 M E}{4\pi} \lim_{P_z \rightarrow \infty} (-i) \int \frac{d^3x}{P_0} \langle P_z | \frac{4}{3} [\partial j_x(\mathbf{x}, t)/\partial t, j_x^\dagger(0)] \\ &\quad - [\partial j_z(\mathbf{x}, t)/\partial t, j_z^\dagger(0)] + \frac{2}{3} i [\partial j_x(\mathbf{x}, t)/\partial t, j_y^\dagger(0)] | P_z \rangle_{t=0}. \end{aligned} \quad (5.8)$$

Therefore, we predict not only that $\bar{\nu}p$ and νp total cross sections depend linearly on energy, but that the sum of the total cross sections is determined by the equal-time commutator of the Cabibbo current with its time derivative at infinite momentum.

The linear rise of cross sections predicted here would be cut off, were there an intermediate boson W exchanged, with the cutoff at $E \sim M_W^2/M_p$. Data from the deep-mine cosmic-ray neutrino experiments^{16,17} are as yet inconclusive; however, a linear rise of neutrino cross sections up to 10–100 BeV is not inconsistent with the data.¹⁸

VI. CONCLUSIONS

By combining the $q_0 \rightarrow i\infty$ asymptotic limit with the infinite-momentum method, we have shown that in a certain limit, the inelastic electron scattering structure functions

$$\lim_{q^2 \rightarrow \infty, -q^2/M\nu = \omega \text{ fixed}} MW_1(\nu, q^2) \equiv F_t(-q^2/M\nu), \quad (6.1)$$

$$\lim_{q^2 \rightarrow \infty, \omega \text{ fixed}} [(-q^2)W_2(\nu, q^2) - W_1(\nu, q^2)] \equiv \frac{F_t(-q^2/M\nu)}{M} \quad (6.2)$$

are directly related to Fourier transforms of almost-equal-time commutators at infinite nucleon momentum,

given in (2.5) and (2.6). Provided the Callan-Gross⁴ integral is finite,

$$\lim_{|q^2| \rightarrow \infty} |q^2| \int_0^\infty \frac{d\nu}{\nu} W_2(\nu, q^2) < \infty, \quad (6.3)$$

we have shown that these commutators are not infinite, but may be zero (or ambiguous). The hypothesis that these commutators are indeed finite is equivalent to the prediction

$$MW_1 \xrightarrow{q^2 \rightarrow \infty} F_1(-q^2/M\nu), \quad (6.4)$$

$$\nu W_2 \xrightarrow{q^2 \rightarrow \infty} F_2(-q^2/M\nu). \quad (6.5)$$

Under similar assumptions, total $\bar{\nu}p$ and νp cross sections are predicted to rise linearly with laboratory neutrino energy. Of particular interest is the behavior of the sum of cross sections, dependent, according to (5.8), only on the equal-time commutator of the Cabibbo current with its time derivative, evaluated between nucleon states at infinite momentum.

An extension of this technique to more general kinematical conditions, presented in Sec. IV, is possible, but by itself does not seem to lead to further insight into the nature of this limit. A more physical approach into what is going on is, without question, needed.

ACKNOWLEDGMENTS

We thank R. Norton, R. P. Feynman, members of the Stanford Linear Accelerator Center (SLAC) theory group, and the SLAC electron-scattering group for useful and enjoyable conversations.

¹⁶ M. Menon *et al.*, Can J. Phys. **46**, S344 (1968).

¹⁷ F. Reines *et al.*, Can J. Phys. **46**, S350 (1968).

¹⁸ For neutrino-scattering from a neutron, the term in (5.8) proportional to $[J_x, J_y^\dagger]$ changes sign, and the others are equal to the corresponding terms for the proton, in the approximation of ignoring $|\Delta S=1|$ transitions. Thus the deep-mine experiments measure mainly the diagonal commutators $[J_z, J_z^\dagger]$ and $[J_x, J_x^\dagger]$ only.