The Pauli objection

Juan Leon¹, Lorenzo Maccone²

¹Instituto de Física Fundamental, CSIC, Serrano 113-B, 28006, Madrid, Spain

²Dip. Fisica and INFN Sez. Pavia, University of Pavia, via Bassi 6, I-27100 Pavia, Italy

Schrödinger's equation says that the Hamiltonian is the generator of time translations. This seems to imply that any reasonable definition of time operator must be conjugate to the Hamiltonian. Then both time and energy must have the same spectrum since conjugate operators are unitarily equivalent. Clearly this is not always true: normal Hamiltonians have lower bounded spectrum and often only have discrete eigenvalues, whereas we typically desire that time can take any real value. Pauli concluded that constructing a general a time operator is impossible (although clearly it can be done in specific cases). Here we show how the Pauli argument fails when one uses an external system (a "clock") to track time, so that time arises as correlations between the system and the clock (conditional probability amplitudes framework). In this case, the time operator is not conjugate to the system Hamiltonian, but its eigenvalues still satisfy the Schrödinger equation for arbitrary Hamiltonians.

There are many different ways in which a time operator [1–3] can be introduced into quantum mechanics. These differences reflect the different physical meanings that "time" may have. We recall the main ones (the following list is, by necessity, incomplete and clearly the following categorizations are not clear-cut): (1) Typical time operators [4–8] represent a "time of arrival", whose measurement represents the time at which a system is in a certain location. This is dual [9] to the Newton-Wigner [10] position operator whose measurement represents the position at which a system is located at a certain time; (2) coordinate time [11–14]; (3) an arbitrary parameter (also reinterpreted as "Newtonian absolute time") [15, 16]; (4) a dynamical variable that parametrizes different Hilbert spaces [17–19]; (5) a classical parameter that cannot be quantized [20-23]; (6) a parameter that can be quantized, but not using self-adjoint operators (observables) [11, 12, 24, 25]; (6) proper time [26]; (7) clock time [27–44].

Here will be dealing with the clock time, mainly focusing on the Page-Wootters and Aharanov-Kaufherr (PWAK) approach [29–33]. In this framework, time is defined as "what is shown on a clock", where a clock is some (external) physical system that is taken as a time reference. Then, the measurement of time acts as a conditioning that outputs the position in time of some event that is being gauged by the clock: namely the emphasis is on the correlations between a system and the clock as in 'the state of a system given that the clock shows t'. As shown in [29–34] these correlations manifest themselves as a "static" entangled global state that satisfies a Wheeler-de Witt [45] equation. The PWAK approach is briefly reviewed in Sec. I.

The Pauli objection [21] essentially states that since the energy is the generator of (continuous) time translations, any time operator must be conjugate to an energy operator (Hamiltonian) that has unbounded continuous spectrum, properties which are not satisfied by the Hamiltonian of typical systems. The standard textbook answer to this objection is that time in quantum mechanics cannot be represented by an operator and is a parameter, external to the theory, e.g. [20, 35]. A different, but connected objection was put forth by A. Peres [46]: if the Hamiltonian is the generator of time translations and the momentum is the generator of space translations, then the Hamiltonian and the momentum must always commute, since space and time are independent degrees of freedom. In this paper we show how the PWAK formalism can easily bypass these objections and provide an acceptable time operator.

The main idea is simple: the global Hamiltonian must contain both the system Hamiltonian (which may have arbitrary spectrum) and the clock Hamiltonian, which for an ideal clock must have unbounded continuous spectrum (physical clocks can clearly only approximate this ideal situation). It is the clock Hamiltonian that is conjugate to the time operator, whereas it commutes with the system Hamiltonian which acts on a different Hilbert space. Then the clock Hamiltonian is the generator of clock shifts, hence of "time" translations, whereas the system Hamiltonian is the generator of translations only of the system state, and not of time. Then [T, H] = 0, so that H and \hat{T} do not need to have the same spectrum. To overcome Peres' objection, one notes that the Hamiltonian indeed does commute with the momentum of the clock.

There are some arguments whose most extreme formulation says that in quantum theory "time is not a quantity at all" [47], i.e., there is no way to attribute values to time, be it an operator or a parameter indistinctly. All these arguments assume that the spectrum of the Hamiltonian generating the time evolution in the state space is bounded from below. Indeed Halvorson ruled out the existence of subspaces of states $s(t_1,t_2)$ that can be associated to time intervals (t_1,t_2) and hence, dispensing with the traditional notion of the passage of time in quantum theory. He concretely derived [47] the contradiction that, as a consequence of the Hegerfeldt theorem [48], $\forall |v\rangle \in s(t_1,t_2) \Rightarrow |v\rangle = 0$. Again, the PWAK approach is immune to these kind of arguments that, as Pauli's, re-

quire boundedness of the Hamiltonian spectrum, which is not the case of the clock Hamiltonian considered here.

While the basic mechanism to overcome the Pauli and Peres objections presented here is clear, one has to be careful, since in the PWAK formalism (reviewed in Sec. I) the dynamics is imposed as a constraint and one must check that even in the space of physical states the above properties still hold true: indeed, in the space of physical states, the Wheeler-de Witt equation "forces" the clock Hamiltonian to coincide with the system Hamiltonian. This analysis is given in Sec. II and III (that contains the more technical parts). Even though the system Hamiltonian and the time operator commute in this framework, it is still possible to give an time-energy uncertainty relation, as shown in Sec. IV.

I. THE PWAK MECHANISM

The PWAK mechanism was initially proposed by Page and Wootters [29–31] and soon after by Aharanov and Kaufherr [32] (but similar previous approaches can be found, e.g. in [34, 37]). A recent review, together with the solution to the objections that were moved against it, can be found in [33].

To provide a quantization of time, one can simply define time as "what is shown on a clock" and then use a quantum system as a clock. If one wants a continuous time that goes from $-\infty$ to $+\infty$, a good candidate clock is to use the position of a 1-d particle [27, 32, 34]. Nonetheless, introducing explicitly a physical system is not necessary, and one can only consider the time Hilbert space as an abstract space with no physical meaning.

The global Hilbert space is then $\mathcal{H}_{TS} = \mathcal{H}_T \otimes \mathcal{H}_S$, where T represents the "time" Hilbert space, typically the one for a particle on a line, $\mathcal{L}^2(\mathbb{R})$. In \mathcal{H}_T we introduce the position operator \hat{T} and the conjugate momentum $\hat{\Omega}$, with $[\hat{T},\hat{\Omega}]=i$. We associate the momentum $\hat{\Omega}$ to the energy of the clock (for a particle, this can be a good approximation for sufficiently massive non relativistic particles [32]). We can enforce that \hat{T} represents the time operator which describes the evolution of a system by imposing the following constraint equation, namely by requiring that the only states $|\Psi\rangle\rangle$ of the joint Hilbert space \mathcal{H}_{TS} that represent physically relevant situations are the ones that satisfy

$$(\hbar\hat{\Omega} \otimes \mathbb{1}_S + \mathbb{1}_T \otimes H)|\Psi\rangle\rangle = 0 , \qquad (1)$$

(*H* being the arbitrary Hamiltonian of the system *S*) which can be interpreted as a Wheeler-de Witt equation [45]. The double ket notation serves only as a reminder that $|\Psi\rangle$ is a state on the joint Hilbert space \mathcal{H}_{TS} . Note that, even though the system Hamiltonian *H* may have arbitrary spectrum, the Wheeler-de Witt Hamiltonian $(\hbar\hat{\Omega} \otimes \mathbb{1}_S + \mathbb{1}_T \otimes H)$ has unbounded continuous spectrum (because $\hat{\Omega}$ has). As eigenstates of the Wheeler-de Witt equation, the physical states $|\Psi\rangle$ are "static" in

the sense that they do not evolve with respect to an "external" time. However, the system evolves with respect to the clock and viceversa, in the sense that the correlations (entanglement) between system and clock track the system evolution. Indeed the solutions of (1) are

$$|\Psi\rangle\rangle = \int_{-\infty}^{+\infty} d\omega \ |\omega\rangle |\tilde{\psi}(\omega)\rangle \ , \tag{2}$$

where $|\omega\rangle$ is the eigenstate of $\hat{\Omega}$ with eigenvalue ω , $|\tilde{\psi}(\omega)\rangle$ is the (un-normalized) Fourier transform of the system state $|\psi(t)\rangle$. [Note that the state (2) is not uniform in $|\omega\rangle$, as the eventual weight is implicit in the norm of $|\tilde{\psi}(\omega)\rangle$, e.g. such weight selects the solutions of (1).] Indeed,

$$|\tilde{\psi}(\omega)\rangle = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dt \ e^{-i\omega t} |\psi(t)\rangle \Rightarrow \tag{3}$$

$$H|\tilde{\psi}(\omega)\rangle = \sqrt{2\pi} \sum_{k} \delta(\omega_{k} + \omega) \psi_{k} \hbar \omega_{k} |e_{k}\rangle = -\hbar \omega |\tilde{\psi}(\omega)\rangle, \tag{4}$$

where we have used the expansion $|\psi(t)\rangle = \sum_k \psi_k e^{-i\omega_k t} |e_k\rangle$ in terms of the Hamiltonian eigenstates $|e_k\rangle$ of eigenvalue $\hbar\omega_k$. If the Hamiltonian has a continuous spectrum, an analogous expression holds:

$$H|\tilde{\psi}(\omega)\rangle = \sqrt{2\pi} \int d\omega' \delta(\omega' + \omega) \psi(\omega') \hbar \omega' |\omega'\rangle$$
$$= -\hbar \omega \sqrt{2\pi} \psi(-\omega) |-\omega\rangle = -\hbar \omega |\tilde{\psi}(\omega)\rangle, \quad (5)$$

where $|\omega\rangle$ is the δ -normalized energy eigenstate of eigenvalue $\hbar\omega$. It is clear from these expressions that $|\tilde{\psi}(\omega)\rangle$ is the null vector if ω is not an eigenvalue of the Hamiltonian. The solutions (2) can be written as

$$|\Psi\rangle\rangle = \int_{-\infty}^{+\infty} d\omega \ |\omega\rangle|\tilde{\psi}(\omega)\rangle = \int_{-\infty}^{+\infty} dt \ |t\rangle|\psi(t)\rangle \ , \quad (6)$$

where $|t\rangle = \int d\omega \, e^{-i\omega t} |\omega\rangle/\sqrt{2\pi}$ is the position eigenstate in \mathcal{H}_T and $|\psi(t)\rangle$ is the system state at time t in \mathcal{H}_S , with normalization $\langle \psi(t)|\psi(t)\rangle = 1$ for all t. [Note that any nontrivial probability amplitude $\phi(\omega)$ in the integral (2) can be absorbed in the definition of the system state $|\tilde{\psi}(\omega)\rangle$ as $\psi_k \to \phi(\omega_k)\psi_k$.] The states (6) are improper (non-normalizable) states that reduce to the momentum eigenstate $\sqrt{2\pi}|\omega=0\rangle = \int dt|t\rangle$ in \mathcal{H}_T whenever the system is in an eigenstate of its Hamiltonian H. Starting from the state (6) and conditioning the clock system to the state $|t\rangle$, we recover the Schrödinger equation: indeed, Eq. (1) in the position representation becomes

$$\langle t | \hbar \hat{\Omega} + H | \Psi \rangle \rangle = 0 \Leftrightarrow (-i\hbar \frac{\partial}{\partial t} + H) | \psi(t) \rangle = 0 \; , \eqno(7)$$

where we wrote the momentum in position representation $\langle t|\hat{\Omega}=(-i\partial/\partial t)\langle t|$, and we used $\langle t|\Psi\rangle\rangle=|\psi(t)\rangle$ which follows from (6). One can similarly also derive the unitary evolution for the system [33].

One of the many advantages of this approach is that it renders explicit the problem that, when an event is gauged by a quantum clock or a system is controlled by a quantum clock, a feedback (disturbance) to the clock must occur [28].

II. BYPASSING THE PAULI OBJECTION

The Pauli objection is just an argument and is not really rigorous. There are many counterexamples in the literature (e.g. [49]), but it can also be made into a rigorous statement if one is careful enough (e.g. [50]). It basically says that if one introduces a time operator, then time and energy are conjugate operators through the Schrödinger equation. Then their spectrum must be the same. This is a consequence of the Stone-von Neumann theorem: if $[\hat{T}, \hat{H}] = i$ then \hat{T} and \hat{H} have the same spectrum.

The PWAK mechanism is immune to this, since we are requesting that $[\hat{T},\hat{\Omega}]=i$ and then enforcing that $\hat{\Omega}$ is equal to \hat{H} only on the physical states through the constraint Eq. (1). Such equation is saying that in this subspace $\hat{\Omega}=\hat{H}!$ So it seems that in the space of physical states, the Pauli argument should apply: \hat{T} has the same spectrum as $\hat{\Omega}$ which (in the subspace) has the same spectrum as \hat{H} . So we must conclude that in the subspace of physical states \hat{T} has the same spectrum as $\hat{H}!$

Luckily this statement is false, although it is not immediately trivial to see. To see why that statement is false, we must formalize it very carefully. We start by defining \hat{T} and $\hat{\Omega}$ as

$$\hat{T} \equiv \int_{-\infty}^{+\infty} dt \ t \ |t\rangle\langle t|, \ \hat{\Omega} \equiv \int_{-\infty}^{+\infty} d\omega \ \omega \ |\omega\rangle\langle\omega| \tag{8}$$

$$|\omega\rangle \; \equiv \; \int_{-\infty}^{+\infty} \frac{dt}{\sqrt{2\pi}} \; e^{-i\omega t} |t\rangle, \; \Rightarrow |t\rangle = \int_{-\infty}^{+\infty} \frac{d\omega}{\sqrt{2\pi}} \; e^{i\omega t} |\omega\rangle.$$

The Pauli objection can be formalized as follows:

- 1. The definitions of \hat{T} and $\hat{\Omega}$ imply that $[\hat{T}, \hat{\Omega}] = i$.
- 2. Introduce the Hilbert space of physical states \mathcal{H}_c as the ones that satisfy Eq. (1), $(\hbar \hat{\Omega} + \hat{H})|\Psi\rangle\rangle = 0$.
- 3. Since $(\hbar\hat{\Omega} + \hat{H})|\Psi\rangle\rangle = 0$, then* also $\hat{T}(\hbar\hat{\Omega} + \hat{H})|\Psi\rangle\rangle = 0$, and $\langle\langle\Phi|\hat{T}(\hbar\hat{\Omega} + \hat{H})|\Psi\rangle\rangle = 0$ for all $|\Phi\rangle\rangle, |\Psi\rangle\rangle \in \mathcal{H}_c$.
- 4. The point above implies that

$$0 = \langle \langle \Phi | \hat{T} (\hbar \hat{\Omega} + \hat{H}) | \Psi \rangle \rangle - \langle \langle \Phi | (\hbar \hat{\Omega} + \hat{H}) \hat{T} | \Psi \rangle \rangle \ (9)$$

$$= \langle \langle \Phi | \hbar[\hat{T}, \hat{\Omega}] + [\hat{T}, \hat{H}] | \Psi \rangle \rangle, \tag{10}$$

so
$$\langle \langle \Phi | \hbar[\hat{T}, \hat{\Omega}] | \Psi \rangle \rangle = - \langle \langle \Phi | [\hat{T}, \hat{H}] | \Psi \rangle \rangle$$
 (11)

5. Since $[\hat{T}, \hat{\Omega}] = i$, this means that, when restricting to the physical states space \mathcal{H}_c , we have $[\hat{T}, \hat{H}] = -i\hbar$, which through the Stone-von Neumann theorem implies that \hat{T} has the same spectrum as \hat{H} in this Hilbert space \mathcal{H}_c , the Pauli objection!

We note that Dirac had introduced an equation of the type (1) in [51], but he did not consider it as a constraint on the physical states. This meant that he ran into an inconsistency similar to the one emphasized above. Dirac never gave a solution [44]. We show here that a solution is provided by the PWAK mechanism.

This above argument is clearly wrong since $[\hat{T}, \hat{H}] = 0$ because they are operators acting on different Hilbert spaces. In fact, the implication indicated with an asterisk at point 3 fails: even though it is true that $(\hbar\hat{\Omega} + \hat{H})|\Psi\rangle\rangle = 0$, this does not imply that $\hat{T}(\hbar\hat{\Omega} + \hat{H})|\Psi\rangle\rangle = 0$. This comes from the fact that the spectrum of \hat{T} is unbounded, see the definition (8). We prove this in the following section (using two different regularizations for the physical states $|\Psi\rangle\rangle$ which are un-normalizable).

One can also give a physical interpretation to this: one should expect that the expectation value of \hat{T} must be undefined in the space of physical states. In fact $\langle \hat{T} \rangle$ has as value the result to the question "what is the time?" which is a meaningless question per se in physics. Meaningful questions are "what is the time when the spin is up?" or "what is the time now that you're reading this?", etc. So, one must expect that $\langle \langle \Phi | \hat{T} | \Psi \rangle \rangle$ will be undefined in the space of physical states, which is indeed what happens.

III. REGULARIZATION

Here we provide the regularizations necessary to prove the relations introduced in Sec. II.

The state (6) is the solution of the eigenvalue equation (1). The eigenvalue $\lambda=0$ is an essential eigenvalue of the self-adjoint constraint operator $\hat{\mathbb{J}}=\hbar\hat{\Omega}+\hat{H}$. This can be shown through Weyl's criterion [52] (Chap. 7), since $\|(\hat{\mathbb{J}}-\lambda)|\Psi_n\rangle\|\to 0$ for $n\to\infty$ where $|\Psi_n\rangle\rangle$ is a Weyl sequence, i.e. a normalized sequence of vectors that weakly converges to zero (namely, $\forall |\theta\rangle\rangle \in \mathcal{H}$ we have $\langle\langle\theta|\Psi_n\rangle\rangle\to 0$).

We will show this using two different Weyl sequences which can be considered as approximate eigenvectors (as expected, both give the same results):

$$|\Psi_n\rangle\rangle \equiv \left(\frac{2}{\pi n}\right)^{1/4} \int dt \ e^{-t^2/n} |t\rangle |\psi(t)\rangle$$
 (12)

$$|\Psi'_{m}\rangle\rangle \equiv \frac{1}{\sqrt{m}} \int dt \, \beta(t/m)|t\rangle|\psi(t)\rangle , \qquad (13)$$

where the first uses a Gaussian whose width diverges for $n \to \infty$, the second uses the box function β whose width diverges for $m \to \infty$, with $\beta(x) = 1$ if $-\frac{1}{2} < x < \frac{1}{2}$, $\beta(-\frac{1}{2}) = \beta(\frac{1}{2}) = \frac{1}{2}$, and $\beta(x) = 0$ otherwise. It has derivative $\partial \beta(x)/\partial x = \delta(x+1/2) - \delta(x-1/2)$. These are both Weyl sequences (see [52] at pages 71 and 74 respectively).

All states in \mathcal{H}_c can be obtained from these as

$$|\Psi\rangle\rangle = \lim_{n \to \infty} \left(\frac{\pi n}{2}\right)^{1/4} |\Psi_n\rangle\rangle = \lim_{m \to \infty} \sqrt{m} |\Psi'_m\rangle\rangle.$$
 (14)

Note that the state $|\Psi\rangle\rangle$ is un-normalizable: it does not live in a Hilbert space, but one has to resort to rigged Hilbert spaces, where the Hilbert space containing normalized vectors (and the limit of sequences of normalized vectors) is incremented with vectors of infinite norm [53].

First we show that $|\Psi_n\rangle\rangle$ and $|\Psi'_m\rangle\rangle$ are indeed Weyl sequences for $\hat{\mathbb{J}}$ and $\lambda=0$, namely they are "proper" approximations of the improper eigenvectors of $\hat{\mathbb{J}}$ with eigenvalue $\lambda=0$. (We already know that, this is just a consistency check.) Let us start with $|\Psi_n\rangle\rangle$. We have to show that $\|(\hat{\mathbb{J}}-\lambda)|\Psi_n\rangle\rangle\|\to 0$ for $n\to\infty$. Indeed,

$$\lim_n \langle \hbar \hat{\Omega} + \hat{H} \rangle |\Psi_n \rangle \rangle = \lim_n \frac{2i}{n} \left(\frac{2}{\pi n} \right)^{1/4} \int dt \ t \ e^{-t^2/n} |t \rangle |\psi(t) \rangle.$$

That this is a null vector can be seen by taking its modulus:

$$\|(\hbar\hat{\Omega} + \hat{H})|\Psi_n\rangle\rangle\|^2 = \frac{4}{n^2}\sqrt{\frac{2}{\pi n}}\int dt \ t^2 \ e^{-2t^2/n} \to 0,$$

where we used the fact that $\int dt \ t^2 \ e^{-at^2} = \sqrt{\pi/a^3}/2$ and $\langle \psi(t)|\psi(t)\rangle = 1$. The same result applies using the other regularization: $(\hbar\hat{\Omega} + \hat{H})|\Psi'_m\rangle\rangle \to 0$ for $m \to \infty$. Indeed, for all vectors $|\theta\rangle\rangle = \int dt \theta(t)|t\rangle|\phi(t)\rangle$ in the Hilbert space, we find

$$\begin{split} & |\langle\langle\theta|(\hbar\hat{\Omega}+\hat{H})|\Psi\rangle\rangle| & (15) \\ &= \lim_{m} |\langle\langle\theta|\int dt \; |t\rangle|\psi(t)\rangle[\delta(t-\frac{m}{2})-\delta(t+\frac{m}{2})] \\ &= \lim_{m} |\langle\phi(0)|\psi(0)\rangle[\theta^*(\frac{m}{2})-\theta^*(-\frac{m}{2})]| = 0 \;, \quad (16) \end{split}$$

where we used (14), and the fact that $\langle \phi(t)|\psi(t)\rangle$ is constant and that all square integrable functions $\theta(t) \to 0$ for $t \to \pm \infty$.

Now the crucial point: what happens when we multiply these null vectors by the unbounded operator \hat{T} ? We obtain a non-null vector! Indeed.

$$\|\hat{T}(\hbar\hat{\Omega} + \hat{H})|\Psi_n\rangle\rangle\|^2 = \frac{4}{n^2}\sqrt{\frac{2}{\pi n}}\int dt \ t^4 \ e^{-2t^2/n} = \frac{3}{4} \ ,$$

for all n, since $\int dt \ t^4 \ e^{-at^2} = 3\sqrt{\pi/a^5}/4$. This implies that $|\Psi_n\rangle\rangle$ is not an approximate eigenvector for the $\lambda=0$ eigenvalue of the operator $\hat{T}(\hbar\hat{\Omega}+\hat{H})$, even though it was an approximate eigenvector for the operator $(\hbar\hat{\Omega}+\hat{H})$. This also means that in the rigged Hilbert space we cannot consider $|\Psi\rangle\rangle$ as eigenvector of this operator.

One can also show that

$$\langle \langle \Psi_n | (\hbar \hat{\Omega} + \hat{H}) \hat{T} | \Psi_n \rangle \rangle$$

$$= i \sqrt{\frac{2}{\pi n}} \int dt (e^{-2t^2/n} - \frac{2t^2}{n} e^{-2t^2/n}) = \frac{i}{2} , \quad (18)$$

which suggests that $|\Psi\rangle\rangle$ is an (improper) eigenstate of $(\hbar\hat{\Omega} + \hat{H})\hat{T}$ with eigenvalue i/2. Indeed, the above results imply

$$\|[(\hbar\hat{\Omega} + \hat{H})\hat{T} - \lambda]|\Psi_n\rangle\rangle\| \to 0 \tag{19}$$

for $\lambda = i/2$ (actually the modulus is equal to 0 for all n). Note that this is the value that is necessary in Eq. (9) to avoid the contradiction!

Analogous considerations hold for the other regularization since

$$\langle \langle \theta | \hat{T}(\hbar \hat{\Omega} + \hat{H}) | \Psi \rangle \rangle = \lim_{m} i \frac{m}{2} [\theta^*(\frac{m}{2}) + \theta^*(-\frac{m}{2})], \quad (20)$$

where we used $|\Psi\rangle\rangle = \lim_m \sqrt{m} |\Psi'_m\rangle\rangle$. This does not tend to zero as $m \to \infty$ for all square integrable functions $\theta(t)$, since square integrable functions must go to zero faster than $1/\sqrt{t}$ for $t \to \infty$.

In conclusion, not only we have shown that point 3 of the Pauli argument fails, but we have also recovered the expected values of the scalar products $\langle \langle \Psi | \hat{T} (\hbar \hat{\Omega} + \hat{H}) | \Psi \rangle \rangle = i/2$ that are necessary in Eq. (9) if it has to be consistent with the fact that $[\hat{T}, \hat{\Omega}] = i$.

IV. UNBOUNDED-ENERGY CLOCKS?

The way we the PWAK mechanism bypasses the Pauli objection is by using a clock with an unbounded Hamiltonian equal to its momentum [32]. Clearly this is unphysical and one could object that our resolution is not a resolution after all. However, it is important to notice that all quantum experiments to date have been performed with macroscopic "classical" clocks (except for especially crafted situations [43]). These have energy so large compared to the time uncertainties that can be tracked in practice that their spectrum can be considered unbounded for all practical purposes. Moreover, macroscopic systems get very quickly correlated to astronomical distances (e.g. the motion of one gram of matter on the star Sirius by one meter sensibly influences the particle trajectories in a box of gas on earth on a time-scale of μ s after the transit time [54]) so that a pure-state analysis as performed above will break down unless one is able to track all the correlated degrees of freedom, a practical impossibility.

In this section we study how good is the approximation of considering a clock with unbounded spectrum. We show that if the energy spread is ΔE the time can be measured up to a precision $\Delta t = \hbar/2\Delta E$. This is a direct consequence of the time-energy uncertainty relation [55–57] which says that, if the energy spread is ΔE , then the minimum time interval it takes to evolve to an orthogonal state is $\tau \geqslant \hbar/2\Delta E$. Hence no smaller time interval can be measured with accuracy.

Clearly, a spread in energy is by itself insufficient to obtain a clock: one also needs good time correlation. However, in the absence of energy spread, a clock in the state $|\Psi\rangle\rangle$ of (2) cannot keep time, and with limited energy spread, it can only keep time up to some accuracy since the correlation in time cannot be sufficiently high. For the sake of simplicity we will keep the unbounded Hamiltonian of the clock $\hbar\Omega$, but we will reduce the energy spread by making explicit the spectral

function $\phi(\omega)$ which was absorbed into $|\tilde{\psi}(\omega)\rangle$ in (2) as $|\tilde{\psi}(\omega)\rangle = \phi(\omega)|\tilde{\chi}(\omega)\rangle$, and we take the standard deviation of the probability probability $|\phi(\omega)|^2$ is $\Delta\omega = \Delta E/\hbar$. Since the clock and the system are entangled, this spectral function does not refer exclusively to the clock, but to both the clock and the system. As expected from the time-energy uncertainty relation, a limited-bandwidth spectral function $\phi(\omega)$ will reduce the speed of evolution (time resolution of the global system). Indeed (neglecting multiplicative constants) we have

$$|\Psi\rangle\rangle = \int d\omega \,\phi(\omega)|\omega\rangle|\tilde{\chi}(\omega)\rangle \propto \int dtdt' \,\tilde{\phi}(t-t')|t\rangle|\chi(t')\rangle,$$

where $\tilde{\phi}$ and $|\chi(t)\rangle$ are the Fourier transforms of ϕ and $|\tilde{\chi}(\omega)\rangle$. Even though this seems to be incompatible with Eq. (6), it is not as can be seen by writing

$$|\psi(t)\rangle \propto \int dt' \tilde{\phi}(t-t')|\chi(t')\rangle$$
 (21)

This can be interpreted as if $|\psi(t)\rangle$ is obtained by "averaging" $|\chi(t)\rangle$ over time with a probability amplitude $\tilde{\phi}$. Then the smallest time interval during which $|\psi(t)\rangle$ can vary appreciatively is of the order of $\hbar/\Delta E$, i.e. the inverse of the spread of the probability $|\phi(\omega)|^2$. Indeed

$$\langle \psi(t)|\psi(t')\rangle \propto \int d\tau d\tau' \,\tilde{\phi}(t-\tau)\tilde{\phi}^*(t'-\tau')\langle \chi(\tau')|\chi(\tau')\rangle,$$

whence, even supposing instantaneous change of $|\chi\rangle$, namely $\langle \chi(\tau')|\chi(\tau')\rangle \propto \delta(\tau-\tau')$, we have

$$\langle \psi(t)|\psi(t')\rangle \propto \int d\omega |\phi(\omega)|^2 e^{i\omega(t-t')}.$$
 (22)

If $|\phi(\omega)|^2$ has a spread $\simeq \Delta E/\hbar$, then its Fourier transform will have a spread of the order of $\hbar/\Delta E$. This means that the scalar product $\langle \psi(t)|\psi(t')\rangle$ cannot change appreciatively in a smaller interval, namely the time scale of change of the system state must be larger than $\hbar/\Delta E$, in accordance with the time-energy uncertainty relation.

Thanks to its linearity, the Schrödinger equation holds for the "averaged" $|\psi(t)\rangle$ of Eq. (21), which implies that eventual imperfect correlations between system and clock will not induce a fundamental decoherence effect in this framework. This contrasts to the Gambini et al. framework [41, 42] where imperfect clocks do induce a fundamental decoherence.

V. CONCLUSIONS

In this paper we have shown how one can easily bypass the Pauli and the Peres objections to a quantum operator for time using the conditional probability amplitude framework of Page, Wootters, Aharanov and Kaufherr. Moreover we have detailed how the time-energy uncertainty relation arises in this framework.

Acknowledgments

We acknowledge the FQXi foundation for financial support in the "Physics of what happens" program. JL acknowledges support from MINECO/FEDER Project FIS2015-70856-P and CAM PRICYT Project QUITEMAD+ S2013/ICE-2801 and Giacomo Mauro D'Ariano for the kind hospitality at Pavia University. LM acknowledges very useful feedback from Vittorio Giovannetti and Seth Lloyd.

- [1] K.V. Kuchař, "Time and interpretations of quantum gravity", Proc. 4th Canadian Conference on General Relativity and Relativistic Astrophysics, eds. G. Kunstatter, D. Vincent, and J. Williams (World Scientific, Singapore, 1992), pg. 69-76.
- [2] E. Anderson, "The Problem of Time in Quantum Gravity" in Classical and Quantum Gravity: Theory, Analysis and Applications, Ed. V.R. Frignanni. Nova, New York 2012.
- [3] K.V. Kuchař, in Quantum Gravity 2: a Second Oxford Symposium, ed. C.J. Isham, R. Penrose and D.W. Sciama (Clarendon, Oxford 1981); K.V. Kuchař, in Conceptual Problems of Quantum Gravity, ed. A. Ashtekar and J. Stachel (Birkhäuser, Boston 1991); C.J. Isham, in Integrable Systems, Quantum Groups and Quantum Field Theories, ed. L.A. Ibort and M.A. Rodríguez (Kluwer, Dordrecht 1993); K.V. Kuchař, in The Arguments of Time, ed. J. Butterfield (Oxford University Press, Oxford 1999).
- [4] Y. Aharonov, J. Oppenheim, S. Popescu, B. Reznik, W. G. Unruh, "Measurement of time of arrival in quantum mechanics", Phys. Rev. A 57, 4130 (1998).

- [5] R. Werner, "Arrival time observables in quantum mechanics", Annales de l'I. H. P., section A, 47, 429 (1987).
- [6] B. Mielnik, "The Screen Problem", Found. Phys. 24, 1113 (1994).
- [7] V. Delgado, J. G. Muga, "Arrival time in quantum mechanics", Phys. Rev. A 56, 3425 (1997).
- [8] E.A. Galapon, A. Villanueva, "Quantum first time-of-arrival operators", J. Phys. A: Math. Theor. 41, 455302 (2008).
- [9] J. Leon, "Time-of-arrival formalism for the relativistic particle", J. Phys. A: Math. Gen. 30, 479 (1997).
- [10] T. D. Newton, E. P. Wigner, "Localized States for Elementary Systems", Rev. Mod. Phys. 21, 400 (1949).
- [11] R. Brunetti, K. Fredenhagen, M. Hoge, "Time in quantum physics: From an external parameter to an intrinsic observable", Found. Phys. 40, 1368 (2010); R. Brunetti, K. Fredenhagen, "Time of occurrence observable in quantum mechanics", Phys. Rev. A 66, 044101 (2002).
- [12] V. S. Olkhovsky, E. Recami, "Time as a Quantum Observable", Int. J. Mod. Phys. A 22, 5063 (2007).
- [13] V.S. Olkhovsky, "Time as a Quantum Observable, Canonically Conjugated to Energy, and Foundations of Self-

- Consistent Time Analysis of Quantum Processes", Adv. Math. Phys. **2009**, 859710 (2009).
- [14] T. G. Downes, G. J. Milburn, C. M. Caves, "Optimal Quantum Estimation for Gravitation", arXiv:1108.5220 (2011).
- [15] E.C.G. Stueckelberg, "La signification du temps propre en mécanique ondulatoire", Helv. Phys. Acta 14, 322 (1941); E.C.G. Stueckelberg, "La mécanique du point matériel en théorie des quanta", Helv. Phys. Acta 15, 23 (1942).
- [16] J. R. Fanchi, "Review of invariant time formulations of relativistic quantum theories", Found. Phys. 41, 4 (2011).
- [17] Y. Aharonov, S. Popescu, J. Tollaksen, "Each instant of time a new Universe", in *Quantum Theory: A Two-Time* Success Story (Springer, 2014) pg. 21-36, arXiv:1305.1615 (2013).
- [18] E. Farhi, S. Gutmann, "The functional integral constructed directly from the Hamiltonian", Ann. Phys. 213, 182 (1992).
- [19] J. Cotler, F. Wilczek, "Entangled histories", Physica Scripta T168, 014004 (2016).
- [20] A. Peres, Quantum Theory: Concepts and Methods (Kluwer ac. publ., Dordrecht, 1993).
- [21] W. Pauli, General Principles of Quantum Mechanics (Springer, Berlin Heidelberg, 1980).
- [22] S. L. Braunstein, C. M. Caves, G. J. Milburn, "Generalized Uncertainty Relations: Theory, Examples, and Lorentz Invariance", Annals Phys. 247, 135 (1996).
- [23] V. Giovannetti, S. Lloyd, and L. Maccone, "Quantum limits to dynamical evolution", Phys. Rev. A 67, 052109 (2003).
- [24] A.S. Holevo, Quantum Systems, Channels, Information (de Gruyter Studies in Mathematical Physics).
- [25] A.S. Holevo, "Estimation of shift parameters of a quantum state", Rep. Math. Phys. 13, 379 (1978).
- [26] D. M. Greenberger, "Conceptual Problems Related to Time and Mass in Quantum Theory", arXiv:1011.3709 (2010).
- [27] H. Salecker, E.P. Wigner, "Quantum Limitations of the Measurement of Space-Time Distances", Phys. Rev. 109, 571 (1958).
- [28] A. Peres, "Measurement of time by quantum clocks", Am. J. Phys. 48, 552 (1980).
- [29] D.N. Page and W.K. Wootters, "Evolution without evolution: Dynamics described by stationary observables", Phys. Rev. D, 27, 2885 (1983).
- [30] D.N. Page, "Clock time and entropy", in *Physical Origins of Time Asymmetry*, eds. J.J. Halliwell, et al., (Cambridge Univ. Press, 1993), arXiv:gr-qc/9303020.
- [31] W.K. Wootters, "'Time' replaced by quantum correlations", Int. J. Theor. Phys. 23, 701 (1984).
- [32] Y. Aharonov, T. Kaufherr, "Quantum frames of reference", Phys. Rev. D 30, 368 (1984).
- [33] V.Giovannetti, S.Lloyd, L.Maccone, "Quantum time", Phys. Rev. D, 92, 045033 (2015).
- [34] P. McCord Morse, H. Feshbach, Methods of Theoretical Physics, Part I (McGraw-Hill, 1953), Chap. 2.6.
- [35] H.D. Zeh, "Emergence of classical time from a universal wavefunction", Phys. Lett. A 116, 9 (1986); H.D. Zeh, Time in quantum theory, http://www.rzuser.uni-heidelberg.de/~as3/TimeInQT.pdf

- [36] V. Vedral, "Time, (Inverse) Temperature and Cosmological Inflation as Entanglement", arXiv:1408.6965 (2014).
- [37] T. Banks, "TCP, quantum gravity, the cosmological constant and all that...", Nucl. Phys. B 249, 332 (1985);
 R. Brout, "On the concept of time and the origin of the cosmological temperature", Found. Phys. 17, 603 (1987);
 R. Brout, G. Horwitz, D. Weil, "On the onset of time and temperature in cosmology", Phys. Lett. B 192, 318 (1987);
 R. Brout, "Time and temperature in semi-classical gravity",
 Z. Phys. B 68, 339 (1987).
- [38] C. Rovelli, "Relational Quantum Mechanics", Int. J. of Theor. Phys. 35, 1637 (1996).
- [39] C. Rovelli, "Time in quantum gravity: An hypothesis", Phys. Rev. D 43, 442 (1991).
- [40] C. Rovelli *Quantum Gravity* (2003), obtainable from http://www.cpt.univ-mrs.fr/~rovelli/book.pdf.
- [41] R. Gambini, R.A. Porto, J. Pullin, S. Torterolo, "Conditional probabilities with Dirac observables and the problem of time in quantum gravity", Phys. Rev. D 79, 041501(R) (2009).
- [42] R. Gambini, L. P. Garcia-Pintos, J. Pullin, "An axiomatic formulation of the Montevideo interpretation of quantum mechanics", Studies in the History and Philosophy of Modern Physics, 42, 256 (2011); R. Gambini, J. Pullin, "The Montevideo interpretation of quantum mechanics: frequently asked questions", J. Phys. Conf. Ser. 174:012003, 2009.
- [43] E. Moreva, G. Brida, M. Gramegna, V. Giovannetti, L. Maccone, M. Genovese, "Time from quantum entanglement: an experimental illustration", Phys. Rev. A 89, 052122 (2014).
- [44] J. Hilgevoord, "Time in quantum mechanics: a story of confusion", Studies in History and Philosophy of Modern Physics 36, 29 (2005).
- [45] B.S. DeWitt, "Quantum Theory of Gravity. I. The Canonical Theory". Phys. Rev. 160, 1113 (1967).
- [46] See [20], Eq. (8.90).
- [47] H. Halvorson, "Does quantum theory kill time?", from http://www.princeton.edu/~hhalvors/papers/, (2010).
- [48] G. C. Hegerteldt and S. N. M. Ruijsenaars, "Remarks on causality, localization, and spreading of wave packets", Phys. Rev D 22, 377 (1980).
- [49] P. Busch, M. Grabowski, P.J. Lahti, "Time observables in quantum theory", Phys. Lett. A 191, 357 (1994).
- [50] M.D. Srinivas, R. Vijayalakshmi, "The 'time of occurrence' in quantum mechanics", Pramana 16, 173 (1981).
- [51] P.A.M. Dirac, "Relativity quantum mechanics with an application to Compton scattering". Proc. Royal Soc. (London) A, 111, 405 (1926).
- [52] P.D. Hislop, I.M. Sigal, Introduction to Spectral Theory, Applied Mathematical Sciences 113 (Springer, 1996).
- [53] L.E. Ballentine Quantum Mechanics, a Modern Development (World Scientific, 2014), Sec. 1.4.
- [54] E. Borel, Le Hasard (Alcan, Paris, 1914).
- [55] L. Mandelstam and I. G. Tamm, "The Uncertainty Relation Between Energy and Time in Non-relativistic Quantum Mechanics", J. Phys. USSR 9, 249 (1945).
- [56] K. Bhattacharyya, "Quantum decay and the Mandelstam -Tamm-energy inequality", J. Phys. A 16, 2993 (1983).
- [57] V. Giovannetti, S. Lloyd, and L. Maccone, "Quantum limits to dynamical evolution", Phys. Rev. A 67, 052109 (2003).